

# MAPPING OF SOIL ORGANIC CARBON CONTENT IN THE GLACIER FORELAND OF MIDTRE LOVÉNBREEN, SPITSBERGEN IN NORTHERN NORWAY



Se-Eun Kim<sup>1)</sup>, Ji Young Jung<sup>1)</sup>, Dominique Laffly<sup>2)</sup>, Lennart Nilsen<sup>3)</sup>, Myrtille Moreau<sup>4)</sup>, Yoo Kyung Lee<sup>1)\*</sup>

- 1) Korea Polar Research Institute, Incheon, South Korea, senny10@kopri.re.kr, jyjung@kopri.re.kr, yklee@kopri.re.kr
- 2) Université de Toulouse, Toulouse, France, dominique.laffly@gmail.com
- 3) University of Tromsø, Tromsø, Norway, lennart.nilsen@uit.no
- 4) GEOREX, Pau, France, myrtille.moreau@gmail.com

## INTRODUCTION

Climate change is rapidly occurring in the Arctic region. Glacier retreat leads to formation of subglacial land, and in turn to proglacial ecosystem. In subglacial area, microorganisms and plants begin to be established, and soil development and soil organic carbon (SOC) accumulation are also initiated. Since space can be substituted with time in the glacier foreland, numerous studies have shown that SOC increases as soil age increase, and such results were drawn from line transect approaches (Hodkinson et al., 2003; Smittenberg et al., 2012). However, glacier foreland is not a simple ecosystem, and SOC accumulation is not only influenced by time. Many factors such as microtopography (altitude, slope, exposition, etc.), radiation, snow free days, runoff, etc, could affect the establishment of biota, and thus impact SOC. Therefore, we aimed to investigate SOC distribution in the glacier foreland with a consideration of many environmental parameters and produce a SOC stock map in the Midtre Lovénbreen.

## MATERIAL & METHODS


### Field Sampling

- Sampling points: 93 sites for vegetation & 34 sites for soil sampling in the foreland of Midtre Lovénbreen in July 2014.
- Soil sampling: 0-5, 5-10, 10-20, and 20-30 cm depths.


### Soil analysis

- SOC & TN content : combustion method at 950 °C after removing total inorganic carbon
- Bulk density : Estimated by soil texture using SPAW software (Soil water characteristics Version 6.02.74; Saxton and Rawls, 2006).
- SOC stock : SOC content (Mg C/Mg soil) x bulk density (Mg soil/m<sup>3</sup> soil) x Soil depth (m).

### Remote sensed data




### Statistical Methods



## ACKNOWLEDGEMENT


This study was supported by an "Environmental Change Studies based on the Arctic Dasan Station: in terms of Geology, Atmospheric Science, and Ecology (PE16030)" funded by the Korea Polar Research Institute.

## STUDY AREA




- **Study area:** Glacier foreland of Midtre Lovénbreen, Brøgger Peninsula, Svalbard, Norway (79°N, 12°W)
- **Sampling sites selection:** Stratified sampling with a consideration of X, Y coordinates, runoff, age, slope, and wind from 300 sites of Moreau (2005)

## RESULTS & DISCUSSION




**Fig 1.** Results of multiple correspondence factorial analysis (MCFA)

- The SOC concentration generally tended to increase as soil age increased but did not always coincide with soil age.
- The active runoff sites showed low concentration of SOC. Glacier/snow meltwater wash out previously established vegetation and accumulated SOC.
- With the same soil age, SOC concentrations were different, this could be due to vegetation establishment affected by microtopography, parent material, climate, living organisms, topography, or time.
- Besides above mentioned factors, we are currently scrutinizing the relationship between SOC and other environmental parameters.
- Through understanding the relationship between SOC concentration/stock and environmental parameters, there is a possibility to quantify and predict SOC distribution by observing vegetation distribution and extracting specific environmental factors.



**Fig 2.** Measured soil organic carbon concentration within 0-5 cm depth



**Fig 3.** Modelling the probability of soil organic carbon (SOC) concentration for 0-5 cm depth in the glacier foreland of Midtre Lovénbreen

## REFERENCE

- Smittenberg, R.H. et al., 2012. Climate-sensitive ecosystem carbon dynamics along the soil chronosequence of the Damma glacier forefield, Switzerland. *Global Change Biol.* 18, pp. 1941-1955.
- Hodkinson, I.D., Coulson, S.J., Webb, N.R., 2003. Community assembly along proglacial chronosequences in the high Arctic: Vegetation and soil development in North-West Svalbard. *J. Ecol.* 91, pp651-663.
- Moreau, M. et al., 2005. Analysis of plant colonization on an arctic moraine since the end of the Little Ice Age using remotely sensed data and a Bayesian approach. *Remote Sens. Environ.* 99, pp. 244-253.
- Saxton, K.E., Rawls, W.J., 2006. Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Sci. Soc. Am. J.* 70.