

JGR Earth Surface

RESEARCH ARTICLE

10.1029/2020JF006030

Special Section:

The Arctic: An AGU Joint
Special Collection

Key Points:

- Detailed subbottom profiler and multibeam echosounder data were acquired at the northwestern Chukchi margin
- The new geophysical data grid reveals a three-dimensional geometry and distribution of glaciogenic sediments and geomorphic features
- Seafloor mapping results indicate four major glacial events with variable grounded ice extent and flow direction

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

S. Kim and Y. K. Jin,
skwan@kopri.re.kr;
ykjin@kopri.re.kr

Citation:

Kim, S., Polyak, L., Joe, Y. J., Niessen, F., Kim, H. J., Choi, Y., et al. (2021). Seismostratigraphic and geomorphic evidence for the glacial history of the northwestern Chukchi margin, Arctic Ocean. *Journal of Geophysical Research: Earth Surface*, 126, e2020JF006030. <https://doi.org/10.1029/2020JF006030>

Received 8 DEC 2020

Accepted 31 MAR 2021

Seismostratigraphic and Geomorphic Evidence for the Glacial History of the Northwestern Chukchi Margin, Arctic Ocean

Sookwan Kim¹ , Leonid Polyak² , Young Jin Joe^{1,3}, Frank Niessen⁴, Hyoung Jun Kim¹, Yeonjin Choi^{1,5}, Seung-Goo Kang¹ , Jong Kuk Hong¹ , Seung-Il Nam¹, and Young Keun Jin¹ 

¹Division of Earth Sciences, Division of Glacial Environment Research, Department of Future Technology Convergence, Korea Polar Research Institute, Incheon, Republic of Korea, ²Byrd Polar and Climate Research Center, The Ohio State University, Columbus, OH, USA, ³Department of Earth and Marine Sciences, Jeju National University, Jeju, Republic of Korea, ⁴Alfred Wegener Institute (AWI), Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany, ⁵Department of Ocean Energy and Resources Engineering, Korea Maritime and Ocean University, Busan, Republic of Korea

Abstract High-resolution seafloor mapping provides insights into the dynamics of past ice sheets/ice shelves on high-latitude continental margins. Geological/geophysical studies in the Arctic Ocean suggest widespread Pleistocene ice grounding on the Chukchi–East Siberian continental margin. However, flow directions, timing, and behavior of these ice masses are not yet clear due to insufficient data. We present a combined seismostratigraphic and morphobathymetric analysis of the Chukchi Rise off the northwestern Chukchi margin using the densely acquired subbottom profiler (SBP) and multibeam echosounder (MBES) data. Comparison with deeper airgun seismic records shows that the SBP data cover most of the glaciogenic stratigraphy possibly spanning ca. 0.5–1 Ma. Based on the stratigraphic distribution and geometry of acoustically transparent glaciogenic diamictites, the lateral and vertical extent of southern-sourced grounded ice became smaller over time. The older deposits are abundant as debris lobes on the slope contributing to a large trough mouth fan, whereas younger grounding-zone wedges are found at shallower depths. MBES data show two sets of mega-scale lineations indicating at least two fast ice-streaming events of different ages. Contour-parallel recessional morainic ridges mark a stepwise retreat of the grounded ice margin, likely controlled by rising sea levels during deglaciation(s). The different inferred advance and retreat directions of the southern-sourced ice reflect complex geomorphic settings. The overall picture shows that the Chukchi Rise was an area where different ice streams had complex interactions. In addition to glaciogenic deposits, we identify a number of related or preceding seabed features including mounds, gullies/channels, and sediment waves.

1. Introduction

Understanding the dynamics of marine-based ice sheets is a critical prerequisite for predicting the responses of modern glacial systems to the warming climate and rising sea levels (e.g., DeConto & Pollard, 2016; Howat et al., 2007). This task requires thorough geological and geophysical investigation of high-latitude continental margins and adjacent borderlands for understanding the buildup and decay mechanisms of the past ice sheets (Dowdeswell et al., 2016, and references therein). While comprehensive data have been collected from the Polar North Atlantic margins since the late twentieth century (Elverhøi et al., 1998), the Quaternary glacial history of the Arctic Ocean remained largely speculative until the last two decades. Recent marine geophysical data employing high-resolution seafloor mapping technologies indicate the extensive impact of grounded ice sheets/ice shelves on the continental margins, as well as bathymetric highs in the central Arctic Ocean (Dove et al., 2014; Jakobsson, 1999; Jakobsson et al., 2008, 2014, 2016; Niessen et al., 2013; Polyak et al., 2001). In particular, a wealth of new data has been collected from the Chukchi–East Siberian margin and the adjacent borderland in the western Arctic Ocean since the late 2000s due to the rapid retreat of sea-ice cover (Coakley et al., 2011; Darby et al., 2005; Jokat, 2009; SWERUS Scientific Party, 2016). The acquired data demonstrate a complex pattern of grounded ice impact on the seafloor interpreted as recurring ice streams advancing from the Laurentide, East Siberian, and Chukchi ice centers (Dove et al., 2014; Jakobsson et al., 2005, 2008, 2014; Niessen et al., 2013; Polyak et al., 2007). This picture


indicates a critical role of this region for understanding the overall Arctic glacial history, while the timing, extent, and provenance of glacial events remain poorly known.

In this paper, we focus on the western side of the Chukchi Rise, a geological structure extending north from the Chukchi continental shelf (Figure 1). The Chukchi Rise and adjacent seafloor structures were shown to have multiple glaciogenic bedforms indicative of ice streams potentially originating from the Laurentide, East Siberian, and Chukchi ice-spreading centers (Dove et al., 2014; Jakobsson et al., 2008, 2014; Polyak et al., 2001, 2007) (Figure 1a). The western Chukchi Rise facing the East Siberian margin is a key but poorly constrained area for understanding these glacial interactions. We analyze new, detailed seismostratigraphic and geomorphic data from this area for empirical constraints on past ice-sheet dynamics and related sedimentary processes. Our analysis is based on the combined high-resolution subbottom profiler (SBP) and multibeam echosounder (MBES) data revealing previously unexplored sedimentary structures, stratigraphy, and submarine landforms. These densely collected geophysical data (Figure 1b) allow for tracing the three-dimensional geometry of individual seafloor and subsurface formations, which provide clues for spatiotemporal distribution of the depositional/erosional glaciogenic features and related seabed forming processes (Dowdeswell et al., 2004, 2016; C. H. Eyles & Eyles, 2010; Rebesco et al., 2016). This empirical evidence augments sediment-core stratigraphy from neighboring areas (Joe et al., 2020; Polyak et al., 2007; Schreck et al., 2018) and ice-sheet modeling studies (Colleoni et al., 2016; Gasson et al., 2018), and thus sheds new light on the glacial history of the Chukchi–East Siberian continental margin. The generated data also provide valuable context for potential scientific drilling projects in the western Arctic Ocean.

2. Background

The Chukchi Borderland in the western Arctic Ocean is composed of the north–south trending bathymetric highs and troughs (Hall, 1990) (Figure 1a). The Chukchi Rise is an immediate extension of the Chukchi Shelf separated by a saddle-like depression from the Chukchi Cap to the north (Hegewald & Jokat, 2013; Shaver & Hunkins, 1964). The Chukchi Basin to the west and the Northwind Basin to the east separate the Chukchi Rise from the East Siberian margin and the Northwind Ridge, respectively. The relatively flat top shelf of the Chukchi Rise has ~200 m water depth (mwd) and less than 0.1° of slope gradient measured along a ~100-km north-south transect (Jakobsson et al., 2020). The western side of the Chukchi Rise shelf has a generally steeper inclination (ca. 0.5°–0.7°) from the crest to the shelf break at ~500–600 mwd than the eastern side (ca. 0.3°). The slope gradient from the shelf break to middle slope at ~1,000–1,100 mwd varies from gentler (ca. 2.6°–3.3°) in the southwestern Chukchi Rise to steeper (ca. 4.6°) in the northwestern Chukchi Rise (white numbers in Figure 1b). The midlower slope gradient (from ~1,100 to 2,200 mwd) on the southwestern margin is ~0.6°. A triangular-shaped bathymetric high, named the western spur in this study, protrudes westward from the middle of the Chukchi Rise ~20 km into the Chukchi Basin (Figure 1b). A broad, seaward-dipping (ca. 0.3°) bathymetric depression (Western Bathymetric Trough) associated with the Herald Canyon on the Chukchi Shelf separates the western slope of the Chukchi Rise from the continental margin (Figure 1b). A similar bathymetric feature on the eastern side of the Chukchi Rise has been named the Broad Bathymetric Trough by Dove et al. (2014). Both troughs are characteristically short and wide (Western Bathymetric Trough ~20–60-km long and 150-km wide) unlike the well-developed cross-shelf troughs on the Arctic continental margins (e.g., Bear Island and M'Clure Strait troughs in Batchelor and Dowdeswell [2014]).

Multiple glaciogenic sediment accumulations and submarine landforms, such as mega-scale glacial lineations (MSGLs), moraine ridges, and iceberg ploughmarks, were identified by the recent geophysical surveys on the Chukchi Borderland (Dove et al., 2014; Hegewald & Jokat, 2013; Ilhan & Coakley, 2018; Jakobsson et al., 2005, 2008; Polyak et al., 2001, 2007) and the East Siberian margin (Joe et al., 2020; Niessen et al., 2013). These seafloor features were interpreted to have been formed by grounding of several-hundred-meters-thick ice caps and/or ice shelves that impacted the Chukchi–East Siberian margin during several Quaternary glaciations. The grounded ice masses on the Chukchi Borderland may have originated from the Laurentide Ice Sheet to the east, the East Siberian margin to the west, and the Chukchi Shelf to the south, possibly with the local ice-cap/ice-sheet covering the shallowest areas such as the Chukchi Plateau (Dove et al., 2014; Jakobsson et al., 2005, 2008, 2014; Polyak et al., 2001, 2007) (Figure 1a). The deep-penetrating airgun seismic profiles across the shelf edge of the Chukchi–East Siberian continental



margin including the Chukchi Rise (purple and blue lines in Figure 1b) show that the marine Neogene deposits on the outer shelf and slope are truncated by glaciogenic sedimentary units downlapping on the preglacial strata (Dove et al., 2014; Hegewald & Jokat, 2013; Ilhan & Coakley, 2018; Niessen et al., 2013). These stratigraphic features indicate that large amounts of sediment were eroded by the grounded ice from the shelf and transported to the slope. Based on the stratigraphic position of the resulting regional unconformity within the Plio-Pleistocene deposits, the Chukchi continental margin has a long glacial history (Hegewald & Jokat, 2013), but its timeline, patterns, and mechanisms involved are still poorly understood.

3. Materials and Methods

All of the new geophysical data used in this study were acquired by the 2015–2019 IBRV *Araon* Arctic Expeditions ARA06B/06C, ARA07C, ARA09C, and ARA10C. The survey lines were collected with a spacing of ~1–3 km on the outer shelf and upper slope of the southwestern Chukchi Rise. These data fill a coverage gap between regional geophysical data of the prior expeditions MGL1112 (Coakley et al., 2011) and ARK-XXIII-3 (Jokat, 2009) (Figure 1b). For a broader regional coverage, we also utilized the 2012 ARA03B data from the more northern and eastern parts of the Chukchi Rise (Figure 1b). This paper is based primarily on the SBP and MBES data, with some of the airgun seismic records used for verifying the identification of the major stratigraphic boundaries. The frequency range of SBP data was set to 2.5–7.0 kHz with the ping rates of 1–2 s depending on the water depth and recording window length (400–500 ms). These settings provided a sediment penetration of tens of meters to less than 100 m, depending on the seabed morphology and geological conditions, with a vertical resolution of ~0.5 m or better. During the ARA10C SBP/MBES survey, the airgun seismic data were simultaneously collected using two Sercel Generator-Injector (G.I.) guns (each 355 in³ volume; 250 in³ for generator and 105 in³ for injector) and a 1.5-km-long Sercel Sentinel solid-type streamer (120 channels). The MBES system recorded travel times and amplitudes of reflected signals with a wide beam angle (–65° to +65°). The swath width of MBES data is ~4.3 times the water depth. The MBES bathymetry data were frequently calibrated using the sound velocity profiles of the conductivity–temperature–depth castings. The MBES backscatter intensities were recorded simultaneously with the bathymetry data acquisition. The quality of all data varied with the weather, sea-ice conditions, and ship speed.

The processing procedure applied to the SBP data includes delay-time shifting, resampling, signal enveloping, and spherical divergence correction using Seismic Unix. The coordinates of each ping point number (PN) were extracted from the SBP data header. The airgun seismic data were processed through conventional processing steps, including setup of the geometry, debubble, velocity analysis, multiple attenuation, prestack time migration, common midpoint stack, and seafloor muting using Schlumberger Omega 2017. The processed SBP and airgun seismic data were imported into the seismic data interpretation software SeisWare Geophysics for acoustic facies analysis, horizon picking, seismic correlation, and isopach mapping. The sediment depths and thicknesses were estimated from the two-way travel time using 1,500 m/s sound speed. The resolved stratigraphic boundaries were traced on the SBP and airgun seismic data (Figures 2 and S1–S3). The identified seismostratigraphic units/subunits were gridded to a resolution of 100 m using a minimum curvature interpolation and were presented in the sediment thickness maps (Figures 3 and S4). The new, high-resolution MBES bathymetry and backscatter data were processed using CARIS

Figure 1. (a) Location map of the Chukchi Borderland in the Arctic Ocean (upper right inset) and a bathymetric map of the Chukchi Borderland (IBCAO version 4.0; Jakobsson et al., 2020). Color-coded arrows indicate ice-flow directions inferred in prior studies (Dove et al., 2014; Jakobsson et al., 2008, 2014; Niessen et al., 2013; Polyak et al., 2001); from the East Siberian (black), Chukchi margin (white), and Laurentide source (red); dashed arrows indicate more tentative interpretations. A black fan-shaped rectangle outlines the study area shown in panel (b). (b) Data location map of the western side of the Chukchi Rise subdivided into the northwestern and southwestern margins and the western spur. Multibeam bathymetry data collected by the IBRV *Araon* (bright-colored) are overlaid on the regional bathymetry (dim colored) with 100-m contour intervals. Thick white lines are the key SBP data shown in Figures 4–7. Labeled thick red segments are locations of the SBP data columns (a)–(f) in Figure 2. The black rectangle outlines the detailed survey area shown in Figures 9a and S5a. Other rectangles show the location of additional ARA03B data (green, Figures 9b and 9c) and the study areas from Polyak et al. (2007) on the Northwind Ridge and Joe et al. (2020) on the Arliss Plateau (red). White numbers show the upper slope gradients. Blue and purple lines are regional geophysical data grids of MGL1112 (Coakley et al., 2011; Dove et al., 2014) and ARK-XXIII/3 (Hegewald & Jokat, 2013; Jokat, 2009), respectively. Thick yellow lines crossing the outer shelf to the upper-mid slope are the multichannel seismic profiles of MGL1112 (c) and ARA10C (Figure S3). Dark gray and blue dashed lines are 600-mwd and 350-mwd isolines marking the shelf edge and the limit of iceberg ploughmarks, respectively. WS, Western spur. (c) Seismic profile MGL1112 crossing the Chukchi Shelf and Chukchi Rise shows crustal blocks completely covered by well-developed sedimentary strata. VE, vertical exaggeration. Modified from Ilhan and Coakley (2018); with permission from Elsevier. SBP, subbottom profiler.




Figure 2. (a–f) The SBP data columns summarizing the seismic stratigraphy of the western Chukchi Rise margin. Seismostratigraphic units U4–U1 are defined by the laterally continuous reflectors marked by thick gray solid lines. Depositional bodies (lenses) T4–T1 are the acoustically transparent subunits (facies) within units U4–U1. The location of the SBP columns (a)–(f) is shown in Figure 1b. Vertical axes are two-way travel time (left) and estimated depth in meter at 1,500 m/s sound speed (right). SBP, subbottom profiler.

HIPS&SIPS. The processed MBES results were gridded to a resolution of 20 m and were superimposed on the regional bathymetry grid (IBCAO V4; Jakobsson et al., 2020) using the QGIS software.

4. Results

4.1. Seismostratigraphy

The SBP data show the stratal geometries, internal structures, stacking patterns, and stratigraphic relationships in the uppermost, mostly ~20–40-m-thick deposits. The covered sedimentary succession was divided into four major seismostratigraphic units designated as U4–U1 from the oldest to the youngest (Figures 2 and S4). The unit boundaries were identified by erosional unconformities and/or otherwise laterally sub-continuous, high-amplitude seismic reflections H1–H5 (where H1 is a seafloor) in both of the SBP and accompanying airgun seismic records (Figures 2 and S1–S3). These reflectors have the best expression at the foot of the western Chukchi Rise slope, from where they were traced upslope using the key SBP lines (Figure S1). Although the SBP grid in the northwestern area is relatively sparse, a consistent stacking pattern of the identified units and their mostly continuous boundaries enable the seismostratigraphic interpretation for most of the western Chukchi Rise margin (Figures 2, 3, and S1).

Each major unit is composed of two seismostratigraphic facies (subunits). One is acoustically stratified and shows parallel to subparallel internal configuration, high-to-medium lateral continuity, and low-to-medium amplitude reflections (Figure 2). The stratified subunits, S4–S1, mimic a smooth or undulated geometry of the underlying unit boundaries in most of the study area (Figures 2b, 2c, and 2e). The other facies type is distinguished in lens- or sheet-shaped deposits with a transparent/semitransparent acoustic signature (Figure 2). The lower boundary of the transparent subunits, T4–T1, typically truncates the internal reflections or the underlying units (Figures 2a–2f). These boundaries correspond to the main reflectors between the units, otherwise the upper and lower boundaries of the transparent subunits pinch out and continue as internal reflections (Figures 2b, 2d, and 2f). The constructed isopach maps show the distribution of sediment




Figure 3. Sediment thickness maps of the acoustically stratified subunits S4–S1 (a, c, e, and g) and the acoustically transparent subunits T4–T1 (b, d, f, and h) on the outer shelf and slope of the western Chukchi Rise. The hillshaded multibeam bathymetry data are shown in gray-tones on the regional bathymetry. The estimated sediment thickness was converted from milliseconds to meters at 1,500 m/s sound velocity. The lower boundaries of S4 and T4 on the midlower southwestern slope were partially defined in the SBP data (Figure 4), the thickness of S4/T4 is tentatively plotted in (a) and (b). Dark gray and blue dashed lines are 600-mwd and 350-mwd isolines marking the shelf edge and the limit of iceberg ploughmarks, respectively.




Figure 4. Uninterpreted (a) and interpreted (b) subbottom profile ARA06B-SBP-BB extended into the Chukchi Basin (see Figure 1b for location). The SBP data show stacked glaciogenic debris lobes of acoustically transparent subunits T4 and T3 interbedded with acoustically stratified subunits S4–S1. Vertical axes are two-way travel time (left) and estimated depth in meter at 1,500 m/s sound speed (right). SBP, subbottom profiler.

thickness of the identified stratified subunits S4–S1 and transparent subunits T4–T1 (Figure 3). Below we describe these stratigraphic divisions from bottom to top.

4.1.1. Unresolved Strata

The oldest deposits captured by the SBP records are represented by the stratigraphically unresolved strata with subparallel internal reflections. At the midlower slope of the southwestern Chukchi Rise, the unresolved strata show a stacking pattern similar to the overlying units U4–U1 (PN 5300–6000 in Figure 4 and PN 3000–6000 in Figure 5). In contrast, on the outer shelf of the western Chukchi Rise, the SBP data show that wavy, subparallel internal reflections of stratigraphically unresolved lower strata are truncated by the base of the major units and subunits (PN 3500–4500 in Figure 6 and PN 2000–12000 in Figure 7). These records are similar to the wavy, continuous reflections that are overlain by the thick package of glaciogenic




Figure 5. Uninterpreted (a) and interpreted (b) subbottom profile ARA06C-SBP-AI showing acoustically transparent T4 and T3 lenses within acoustically stratified sediments of S4–S1 on the middle to lower southwestern slope (see Figure 1b for location).

sediments along the slope of the Chukchi margin as shown by the prior and ARA10C airgun seismic data (Dove et al., 2014; Hegewald & Jokat, 2013; Ilhan & Coakley, 2018) (Figures 1 and S3).

4.1.2. Seismostratigraphic Unit U4

The mapped distribution of U4 shows that a thick (>10 m) stratified portion of the major unit, S4, is widely developed along the upper-mid slope of the western spur and the northwestern margin of the Chukchi Rise (white dashed outlines in Figure 3a). The transparent subunit T4 is characterized by ~5-km wide, 10–20-m-thick accumulations along the upper-mid slope of the margin at water depths of 600–1,700 m (Figure 3b). These transparent sediment bodies are intercalated with acoustically stratified sediments (Figure 3b and PN 4800–5800 in Figure 5). Defining the full thickness of the S4 and T4 deposits on the midlower southwestern slope is mostly not possible because most of the lower unit boundaries here are not reached by the SBP data (Figure 2d and PN 2300–5300 in Figure 4). Based on the partially observed lower boundaries of S4 and T4 (PN 1800–2300 and 5400–5500 in Figure 4), we determine the thickness of the S4 and T4 deposits as <10 and ~20–70 m, respectively (Figures 3a and 3b). The upper part of U4 in this area shows a distinct lens-shaped, transparent deposit of T4 (Figures 2d and 4). The overall distribution of U4 is restricted




Figure 6. Uninterpreted (a) and interpreted (b) subbottom profile ARA09C-SBP-AL showing acoustically transparent deposits T2 and T3 interbedded with acoustically stratified sediments on the southwestern outer shelf to upper slope (see Figure 1b for location).

to water depths deeper than 600 m (black dashed line in Figures 3a, 3b, and S4a), being possibly eroded at shallower depths, where younger deposits of U2/U1 rest directly on the erosional surface of the unresolved older strata (Figures 6 and 7).

4.1.3. Seismostratigraphic Unit U3

The stratified subunit S3 is mostly observed on the continental slope of the northwestern margin and the western spur (Figure 3c), whereas the transparent subunit T3 occupies the southwestern margin at water depths of 550–2,100 m (Figure 3d). A 10–20-m-thick accumulation of S3 (white dashed outline in Figure 3c) partially intercalated with transparent lenses of T3 occurs beyond the present-day shelf break (Figure 3d and PN 4800–5000 in Figure 5), similar to the underlying U4 deposits. T3 shows the highest accumulation (>30-m thick, ~70-km wide, white dashed outline in Figure 3d) on the lower slope of the southwestern margin at water depths of 1,800–2,100 m (Figures 2d and 4). This depocenter can be divided into two depositional bodies interbedded with stratified sediments (Figures 2d and 4b). The upper subunit T3a shows




Figure 7. Uninterpreted (a) and interpreted (b) shelf-edge parallel subbottom profile ARA09C-SBP-CT at the northwestern shelf (see Figure 1b for location). The surface of the acoustically transparent deposit T1 features proglacial or subglacial channels and iceberg ploughmarks at water depths below and above 350 m, respectively. (c) MBES backscatter intensity on the top of T1. Red to blue colors show higher to lower backscatter intensity, respectively. MBES, multibeam echosounder.

a lens-shaped external geometry and a relatively restricted distribution, whereas the lower subunit T3b is larger both laterally and vertically and has irregular or sheet-like external geometry (Figure 4). This distinctive stacking pattern of T3 can be observed from the upper to the lower slope of the southwestern Chukchi Rise. T3b is separated from irregularly shaped T4 deposits by a discontinuous, weak reflection (PN 2000–5000 in Figure 4). This faint reflection correlates with the U4/U3 unit boundary in the Chukchi Basin (PN 2000 in Figure 4) and on the lower slope of the western spur (PN 5500 in Figure 4), where acoustic facies are well stratified. On the southwestern outer shelf and upper slope, T3 has a high-amplitude, subcontinuous upper boundary, and a less distinct lower boundary (PN 500–2000 in Figure 6).

4.1.4. Seismostratigraphic Unit U2

A thick (>10 m) stratified subunit S2 is widely developed along the slope of the northwestern margin and the western spur and extends to the southwestern margin of the Chukchi Rise (white dashed outline in Figure 3e). S2 thins out in a basinward direction and reaches a thickness of <10 m in the lower slope (Figure 3e). The internal S2 reflections are generally subparallel to the lower unit boundary of U2 on the midlower slope of the southwestern margin and most of the slope at the northwestern margin and the spur (Figure 5).

The main distribution of T2 shows a >20-m-thick, elongated sediment wedge (Figure 6) extending to >80 km along the shelf edge at water depths of 520–640 m (white dashed outline at the shelf edge in Figure 3f) with a downslope extension of debris lobes to 800 mwd (Figure 8a). The wedge at the shelf edge has an uneven lower boundary truncating the stratified internal reflections of the underlying sediments (PN 1700–3000 in Figure 6). The wedge surface is irregular, and the boundary with the overlying stratified sediments of S2/S1 is distinct on the upper slope and shelf edge but gets disturbed further shelfward (PN 4200–5000 in Figures 6 and 8b). Another >20-m-thick, elongated wedge of a similar deposit occurs further shelfward at 300–450 m depths (white dashed outline at the inner shelf in Figure 3f). The stratigraphic relationship between these adjacent deposits is not completely clear, but the presence of an inclined internal reflector (PN 3300–4000 in Figure 8c) indicates that the wedges probably belong to separate depositional events.

A spatially smaller but comparably thick depocenter attributed to T2 is observed on the middle southwestern slope at water depths of 1,200–1,450 m (Figure 3f). Minor portions of T2 are also distributed on the outer shelf and upper slope of the western spur and the northwestern margin (Figure 3f).

4.1.5. Seismostratigraphic Unit U1

The uppermost seismostratigraphic unit U1 features an overall thin (<10 m) stratified subunit S1 along most of the western Chukchi Rise margin (Figure 3g). U1 thins from 5 to 10 m on the outer shelf to just a couple of meters on the lower slope of the western margin (Figures 2a, 2c, 3g, and S4d). On the outer northwestern shelf, U1 directly overlies the eroded unresolved strata (Figure 7). Transparent facies of T1 cannot be identified in the thin U1 deposits on the midlower slope and rise (Figures 3h, 4, and 5), but show a >20-m-thick and >40-km-long sediment wedge at the northwestern shelf edge at water depths of 310–420 m (Figure 3h). This deposit overlies the unresolved strata with an uneven, erosional unconformity, whereas its surface features high-amplitude incisions and furrows, and is covered by a thin (<1–2 m), conformable sedimentary layer (PN 6200–13000 in Figure 7). Minor amounts of transparent deposits occurring on top of the inner T2 wedge on the southwestern margin likely also belong to T1 (PN 2600–4000 in Figure 8c). Thin S1 deposits are observed at <350-mwd throughout the entire study area (PN 1850–2700 and 11000–13000 in Figure 7). The MBES backscatter data on the U1 surface show a sharp intensity variation from –34 to –28 dB on the unresolved, old strata to –42 to –38 dB on the subunit T1 (Figure 7c).

4.2. Geomorphic Features

The MBES bathymetry data provide three-dimensional geometry and detailed morphological characteristics of submarine landforms related to various subglacial, ice-marginal, glaciomarine, and open marine environments (Figures 9 and S5). In addition, the MBES backscatter data can provide indirect characteristics of seabed such as grain-size composition, sediment compaction, and seafloor roughness and hardness (Fransner et al., 2018; Huang et al., 2018; Todd et al., 2007). As the hull-mounted MBES systems have a lower frequency range (12 kHz) than usual side-scan sonar systems (550 kHz to 1 MHz), the MBES backscatter data in this study are more suitable for detecting differences in bulk density between young, soft muds and older, consolidated sediments rather than grain-size effects (Wille, 2005; Zaragosi et al., 2000).

Two sets of streamlined linear landforms are observed in the MBES bathymetry data on the outer shelf and middle slope of the western spur (Figures 8d and 8e, and sky blue lines in Figure 9a). The S–N trending, 5–12-km-long and 100–200-m-wide lineations on the outer shelf are distributed across a 6-km-wide field at water depths of ~350–460 m (Figures 8d and 9a). At the 350-mwd limit, lineations are overprinted by the randomly oriented, curvilinear to sinuous furrows 50–250-m-wide and <10-m deep (short black lines in




Figure 8. SBP data on the outer shelf to lower slope of the western Chukchi Rise (see Figure 9a for location). (a) Gullies on top of the transparent subunit T2 on the upper southwestern slope. (b) Mounds and hummocky moraines on the outer southwestern shelf. (c) T2 deposits on the southwestern shelf. A faint inclined internal reflector separates two grounding-zone wedges (Figure 3f). (d) Morainic ridges and mega-scale glacial lineations (MSGLs) on the outer shelf to upper slope of the western spur. A cross profile (red solid line) shows the morphology of the MSGL. (e) Buried MSGLs on the middle slope of the western spur. Distinct >10 m high undulations of the seafloor are shown in the cross profile. (f) Buried gullies on the northwestern upper-mid slope. Simultaneously, acquired airgun seismic data are shown in Figure S2. (g) Buried sediment waves on the middle slope of the western spur. SBP, subbottom profiler.

Figure 9a). Both the lineated and furrowed areas have a rough relief appearing in the SBP data as diffraction hyperbolas (e.g., PN 11000–13000 in Figure 7a and PN 7200–8700 in Figure 8d). These areas also feature relatively high MBES backscatter of -32 to -28 dB (red-colored field in Figure 9a) and only very thin sediment cover on top (PN 7200–8700 in Figure 8d). The other lineation set features SSW–NNE trending linear features 4–7-km long and 180–400-m-wide at water depths of 780–940 m on the middle slope (PN 19100–20300 in Figures 8e and 9a). These lineations are formed on top of the transparent subunits T3/T4 and are overlain by 10–15-m-thick acoustically stratified sediments S1/S2 with low backscatter intensities of -42 to -34 dB (Figures 8e and 9a).

Mostly bathymetry-parallel, nested, curvilinear to sinuous ridges are observed along the shelf edge between 410 and 550 mwd of the western spur and the southwestern Chukchi Rise margin (Figure 8d and blue lines




Figure 9. (a) Interpreted submarine landforms drawn on the MBES backscatter map overlapping the hillshaded MBES bathymetry map (see Figure S5 for uninterpreted MBES bathymetry image). The interpreted morphological features, which were formed under subglacial, ice-marginal, and glacialine subenvironments, are marked by differently colored lines and labeled for the explanation. Insets with the SBP image fragments from Figure 8 show examples of acoustic facies corresponding to the submarine landforms. An inset in the lower right corner shows an enlarged view of the hummocky terrain (black box). Labeled white lines show the location of the SBP data in Figures 8a–8g. Ancillary MBES bathymetry data (ARA03B) from the northern (b) and eastern (c) parts of the Chukchi Rise (see Figure 1b for location). MBES, multibeam echosounder; SBP, subbottom profiler.

in Figure 9a). Potentially similar, but shorter and sparser features occur on the outer shelf of the southwestern margin, in the area of a >3-km-wide bathymetric terrace (Figure 9a). The ridges on the western spur are 3–6-m high and >50-km long along the shelf edge (Figure 9a). The deepest ridge on the western spur shows an asymmetric geometry and is covered by a ~3-m-thick stratified sediments of S1 (inset in Figure 8d). The sparser ridges on the southwestern outer shelf are ~3–5-m high and ~3–7-km long (Figure 9a). They are positioned on top of the undulated surface of T2 and are overlain by the stratified sediments of S1 (PN 2200–2700 in Figure 6).

Mound-shaped bedforms with transparent acoustic facies are observed on the upper slope and the outer-shelf terrace of the southwestern margin at water depths of 600–700-m (inside white dashed outlines in Figure 9a). The mounds are aligned subparallel to the boundary between the western spur and the shelf-edge terrace (Figure 9a). Eight mounds surveyed on the upper slope have widths of 200–700 m and are ~10 m higher than the surrounding seafloor (Kim et al., 2020). Larger scale mounds observed on the shelf-edge terrace extend stratigraphically to the U2/U1 unit boundary (PN 2100–2500 in Figures 8b and 9a).

A field of slightly elongated to nearly circular small mounds (hummocks) is located ~5 km shelfward from the shelf break at the southwestern margin at water depths of 440–540 m (inside orange-dashed outline and lower right inset in Figure 9a). This hummocky terrain shows gentle undulations (<10 m high) and a low-to-moderate linearity. The sediments composing these bedforms belong to the transparent subunit T2 covered by stratified sediments of S1 (PN 3000–4500 in Figure 8b).

Narrow, ~100 m wide, depressions (incisions) in a dip direction are observed along the western Chukchi Rise margin beyond the present-day shelf break at water depths of 600–1,600 m (red lines in Figure 9a). The incisions on the southwestern margin are developed on top of the transparent subunit T2 and are covered by ~6-m-thick sedimentary layer S1 (Figure 8a). The depths of these features decrease from ~10–20 m on the shelf break to <10 m on the midlower slope. The upper slope incisions disappear on the western spur

and are observed again on the northwestern margin (Figure 9a). The latter features of ~20–30-m deep and ~400–500-m wide show a more apparent, less sinuous geometry than on the southwestern slope. A thick (~50 m) acoustically stratified sediment with diffraction hyperbolae covers the area with these incisions (PN 700–800 and 1500–2400 in Figures 8f and S2). Some dip incisions ~7–15-m deep and ~500–1000-m wide were found on the outer shelf of the northwestern margin at 370–400-mwd by the narrow stripes of the MBES data (Figure 7c). The SBP data show that these incisions are developed on top of the transparent subunit T1 (PN 6200–11000 in Figure 7).

Distinctive undulated seafloor features elongated parallel to the bathymetry are observed in the unresolved strata on the middle slope (1,300–1,600 mwd) at the western spur and the northwestern Chukchi Rise margin (yellow lines in Figure 9a). On the northern slope, these undulated features are cut by downslope incisions (Figures 8(f) and 9(a)). A thick (>20 m) stratified sedimentary succession of U4–U1 conformably overlies the undulating surface (Figure 8g).

5. Discussion

5.1. Origin of Identified Sedimentary/Geomorphic Features

5.1.1. Seismostratigraphic Units and Facies

The major seismostratigraphic unit boundaries usually indicate either pronounced change in sedimentary environments or depositional unconformities (Veeken & van Moerkerken, 2013). On the glaciated shelves, including the study region (e.g., Dove et al., 2014; Hegewald & Jokat, 2013; Niessen et al., 2013), these features were shown to be characteristically related to the ice-sheet dynamics and associated sedimentary/geomorphic processes. In addition to the direct erosion and deposition by grounded ice, laterally continuous, high-amplitude reflectors commonly occur in glaciomarine environments in connection with massive deposition of iceberg-rafted debris (IRD) during glacial events (Gulick et al., 2017; Joe et al., 2020). The detrital carbonate IRD layers are spatially distributed basin-wide in the western Arctic Ocean and show significantly higher *P* wave velocity and density than the underlying and overlying fine-grained muds (Matthiessen et al., 2010; Schreck et al., 2018; Stein et al., 2010). In particular, a sediment core–seismic correlation in the western part of the Chukchi Basin indicates that strong reflectors in the SBP data correspond to the prominent, IRD-rich, detrital carbonate layers related to major pulses of iceberg discharge from the Laurentide Ice Sheet (Joe et al., 2020).

Seismostratigraphic units/subunits encompassed by the resolved boundaries can be distinguished by the acoustic signature, such as stratified versus transparent facies (Figure 2). According to previous lithostratigraphic studies, the sediments younger than midlate Pleistocene in the Chukchi-East Siberian margin outside the glacially impacted areas mainly consist of fine-grained to sandy muds with cyclic interlamination of brown and yellow to grayish intervals (e.g., Joe et al., 2020; Matthiessen et al., 2010; Park et al., 2020; Schreck et al., 2018; Stein et al., 2010; Xiao et al., 2020). This stratigraphy has been correlated with acoustically stratified facies similar to subunits S4–S1 in areas adjacent to this study area (Dove et al., 2014; Joe et al., 2020; Polyak et al., 2007). A detailed sedimentologic analysis in the western part of the Chukchi Basin indicated that the acoustically stratified facies with the subparallel internal reflectors correspond to poorly sorted sandy muds containing randomly scattered coarse particles affected by drifting iceberg/sea ice (Joe et al., 2020). The lateral continuity and amplitude of the internal reflectors is relatively low compared to the distinctive major unit boundaries (Figure 5). These acoustic characteristics can be explained by the lower concentration of IRD (Joe et al., 2020) possibly due to less intensive ice rafting. Sedimentary layers with an almost constant thickness between the subparallel internal reflectors were interpreted by fine-grained muds deposited by hemipelagic settling or suspension settling of turbid meltwater plumes (Joe et al., 2020), indicating uniform sedimentation conditions (Veeken & van Moerkerken, 2013). The interpretation is consistent with multiple studies of ice-distal glaciomarine sediments (e.g., C. H. Eyles et al., 1991; Ó Cofaigh et al., 2003).

The transparent deposits (e.g., subunits T4–T1 in Figure 2) indicate lenses of acoustically homogeneous sediments without internal bedding due to fast deposition or internal deformation, which is especially common under proglacial or subglacial conditions (Alley et al., 1986; Dowdeswell et al., 2004; Gulick et al., 2017). Resulting deposits are seen in the seismic profiles as sheet- or wedge-shaped, acoustically transparent bodies

usually interpreted as deformation tills (e.g., Dove et al., 2014; Ó Cofaigh et al., 2005). A similar acoustic signature may also characterize deposits formed by episodic events like ice-sheet bulldozing, debris flows, submarine landslides, or subglacial underwater flows during the ice-sheet advances or early deglaciations (e.g., Donda et al., 2008; C. H. Eyles & Eyles, 2010; Gales et al., 2016; Joe et al., 2020; Ó Cofaigh et al., 2003; Vorren & Laberg, 1997). However, these short-lived events can hardly account for major, spatially continuous units.

Overall, sediment depocenters and stratal geometries of the acoustically transparent and stratified sediments (Figures 3–7 and 10a) characterize glaciogenic sedimentary processes and provide useful insights into past ice-sheet dynamics. The seismic stratigraphy of the transparent deposits T4–T2 identified at the southwestern margin (Figures 4–7) corresponds to the prograding foreset beds with an oblique tangential geometry reported for this area based on the airgun seismic data (Dove et al., 2014; Hegewald & Jokat, 2013; Ilhan & Coakley, 2018) (Figures 1c and S3). The outer shelf to slope location, erosional bottom boundary, wedge-shaped stacking pattern, and moderate lateral continuity of these foresets are consistent with seismostratigraphic characteristics of glacial trough mouth fans (TMFs), diamicton-dominated sediment accumulations typically formed at the cross-shelf glacial trough mouth (Batchelor & Dowdeswell, 2014; Dowdeswell & Siegert, 1999; Dowdeswell et al., 1996; Laberg et al., 2000; Ó Cofaigh et al., 2003; Vorren & Laberg, 1997). The occurrence of TMFs at the southwestern Chukchi Rise margin indicates a large amount of sediment transferred by ice sheets to the shelf edge and further down the slope over several glacial cycles. However, in contrast to the typical bulging TMFs at the mouth of High Arctic cross-shelf troughs (Batchelor & Dowdeswell, 2014), the bathymetric contours of the outer shelf and upper slope at the southwestern Chukchi Rise mostly remain slope-parallel or curve slightly inward (Figures 1b and 9a). This morphology can be explained by relatively low rates of sediment supply and/or more available accommodation space (Montelli et al., 2018; Ó Cofaigh et al., 2003). Unstable, short-lived marine-based ice sheets inferred for the Chukchi–East Siberian margin (Batchelor et al., 2019, and references therein) may not be favorable for supplying enough sediment to form an outward bulging slope fan.

The location of the T4/T3 lens- or sheet-shaped depocenters on the midlower slope in the TMF area (Figure 10a) suggests that they have been primarily formed by the glacially fed debris flows, whereas intercalated stratified facies likely indicate suspension settling from turbid meltwater plumes (Joe et al., 2020; Matthiessen et al., 2010; Ó Cofaigh et al., 2003) (Figures 4 and 5). A similar picture has been reported for the East Siberian slope across the Chukchi Basin from the study area (Joe et al., 2020; Niessen et al., 2013). Two T3 subunits separated by the stratified sediments (Figure 4) indicate two depositional events probably reflecting two glacial advances or deglacial pulses.

The T2 depocenter with an erosional basal boundary observed at the outer shelf to upper slope of the Chukchi Rise is interpreted as a grounding-zone wedge (Figures 6 and 10a). Similar deposits composed of diamictons are typically formed at the stable grounding zones of ice sheets with subglacial sediment transportation and deformation (e.g., Batchelor & Dowdeswell, 2015; Batchelor et al., 2018; Dowdeswell & Fugelli, 2012; Newton et al., 2017; Ottesen & Dowdeswell, 2009). Based on the position of the large T2 grounding-zone wedge, during the peak glaciation, the grounding zone was stretched along the shelf break at the modern depths of 550–580 mwd (Figure 6; Figure 5 in Dove et al., 2014). A smaller grounding-zone wedge at depths of ~300–450 m on the southwestern inner shelf (Figures 8c and 10a) possibly indicates another glacial event. The existing data, however, cannot resolve whether this deposit corresponds to T2 or a younger transparent subunit T1. The distribution of the T1 depocenter further north is limited to similar water depths not exceeding ~450 m (Figures 7 and 10a), which suggests that this glaciogenic feature may correspond to the same glaciation. A more detailed characterization of this deposit is not yet available due to a sparse data coverage at the northwestern margin (Figure 1b), notably lacking the shelf-to-slope lines crossing the depocenter.

5.1.2. Submarine Landforms

The randomly distributed, curvilinear to sinuous furrows abundant at water depths shallower than ~350 m (Figures 9a and 9c) are widely interpreted as iceberg ploughmarks generated by the keels of free-floating icebergs under the glaciomarine environment (e.g., Batchelor et al., 2018, and references therein). These features are ubiquitous at <~350-m depths at the outer continental margin from the Canadian Archipelago




Figure 10. (a) Spatial distribution of the acoustically transparent deposits T4–T1 thicker than 20 m. White arrows show major ice-flow trajectories inferred from the distribution of glaciogenic deposits T4–T1 and related geomorphic features. The black rectangle indicates panel (b). (b) Distribution of MSGL (purple lines with yellow arrows for the inferred direction), recessional morainic ridges (blue lines), and proglacial gullies (red lines). The orange-dashed line indicates the inferred maximum extent of the grounded ice. Blue, sky blue, and black dashed lines are projected grounding lines at the present shelf break, 550 mwd, and 500 mwd, respectively. See also schematic cross sections for grounding lines projections in Figure S6. Black arrows indicate retreats of grounding line from the shelf break. MSGL, mega-scale glacial lineation.

to East Siberia and the adjacent borderlands (e.g., Dove et al., 2014; Hunkins et al., 1962; Jakobsson et al., 2005, 2008, 2014; Polyak et al., 2001, 2007).

The long, streamlined landforms mapped in two areas of the western Chukchi Rise margin and the ancillary data (Figures 9a and 9c) are identified as MSGSLs, consistent with the interpretation in multiple studies from the glaciated margins (Dowdeswell et al., 2004; Ottesen et al., 2005; Spagnolo et al., 2014). MSGSL sets have been reported from multiple sites across the western Arctic Ocean margins and borderlands (Dove et al., 2014; Engels et al., 2008; Jakobsson et al., 2008, 2014, 2016; Niessen et al., 2013; Polyak et al., 2001, 2007). In general, these distinct directional features track the fast-flowing ice streams, typically triggered by flotation and abrupt retreat of the grounded ice with rapid deglacial sea-level rise (e.g., Alley et al., 2007; Batchelor et al., 2018; Dowdeswell et al., 2008) (yellow arrows in Figure 10b). Both MSGSL sets in our data occur on a relatively elevated topography, with their general orientation subparallel to the local shelf edge (Figures 9a and 10b), unlike typical MSGSL in the cross-shelf glacial troughs (e.g., Batchelor & Dowdeswell, 2014; Dowdeswell et al., 2008; Halberstadt et al., 2016; Shipp et al., 1999). This pattern suggests that the MSGSL were probably formed by relatively short-lived, passing ice flow from a thicker ice sheet, potentially formed on the continental margin. The episodic nature of such an ice flow may have been beneficial for the MSGSL preservation. We cannot exclude also that these bedforms could be “plane furrows” produced by huge, tilted tabular icebergs derived from a collapsing ice shelf (Dowdeswell & Bamber, 2007; Wellner et al., 2006). Such furrows are relatively sparse in comparison with the MSGSL fields but have a higher linearity than typical iceberg ploughmarks such as observed on the Chukchi margin at water depths shallower than ~350 m (Figure 9). It is not uncommon that MSGSL and “plane furrows” cooccur in areas of the past ice-flow/ice-shelf breakup (Wellner et al., 2006). Another explanation for the MSGSL-like bedforms involves multiple deep-keeled icebergs carried by a surface current along with sea ice, such as on the Yermak Plateau at the exit from the Arctic Ocean (Gebhardt et al., 2011). However, this scenario is unlikely applicable to the Chukchi Basin, where circulation, and thus ice drift is much more sluggish and has a more gyre-like trajectory both in modern conditions and glacial simulations (Stärz et al., 2012).

By comparison with other glaciated seafloor areas, the bathymetry-parallel ridges at the shelf edge (Figure 9) can be interpreted as recessional moraines formed by the stepwise retreat of a grounded ice margin (Batchelor et al., 2017, 2018; Dove et al., 2014; Niessen et al., 2013; Ottesen & Dowdeswell, 2006; Polyak et al., 1997). The acoustically transparent facies of the identified ridges (inset in Figure 8d) indicate poorly sorted deposits, consistent with the diamict material typical for similar features (Bennett et al., 1996; Bradwell et al., 2008).

Mound-shaped morphological features can be formed by the upward migration of gas/fluids, such as free gas from beneath the gas-hydrate stability zone (e.g., Paull et al., 2008; Waage et al., 2019). This interpretation is supported by geochemical data from the mounds mapped on the southwestern upper slope (Figures 8b and 9a) indicating gas hydrates and CH₄ sourced from the deep sedimentary strata (Kim et al., 2020). Such processes can be generated or reactivated by pressure and temperature changes in the deep strata caused by glacial isostatic rebounds during the grounded ice-sheet advances and retreats (e.g., Himmler et al., 2019).

The field of small mounds on the southwestern outer shelf has geomorphic and stratigraphic characteristics similar to hummocky moraines (Elvenes & Dowdeswell, 2016; Ottesen & Dowdeswell, 2009) (Figures 8b and 9a). Such landforms can be formed by stagnant ice pressing onto the soft deformable bed over large areas of submarginal and proximal proglacial areas, including submarine environments (e.g., Batchelor et al., 2018, and references therein; Boone & Eyles, 2001; N. Eyles et al., 1999).

The narrow dip incisions at the shelf edge and on the upper-mid slope (Figures 8a, 8f, and 9a) can be interpreted as gullies, which are commonly developed in front of the past ice grounding lines (e.g., Engels et al., 2008; Gales, 2013; Rydningen et al., 2015). On the polar continental margin, slope gullies can result from sustained cascading of cold waters (brines) and/or sediment-laden subglacial meltwater flows (e.g., Anderson, 1999; Ivanov et al., 2004; A. Lowe & Anderson, 2002; Noormets et al., 2009; Ó Cofaigh et al., 2018; Pope et al., 2018). Broader incisions on top of the youngest transparent deposit T1 on the outer shelf (Figure 7c) may represent valleys formed by the proglacial or subglacial drainage during deglaciation (e.g., Stewart & Lonergan, 2011).

The regular undulations elongated along the slope in preglacial deposits (Figures 8g and 9a) appear like sediment waves formed by persistent along-slope bottom current (e.g., Miramontes et al., 2016). Similar features can also be identified in the airgun seismic records from the Chukchi margin slopes in the preglacial deposits above the upper Miocene unconformity (Hegewald, 2012; Hegewald & Jokat, 2013). A sandy composition of preglacial sediments recovered on the northern Northwind Ridge corroborates a strong current activity at the intermediate water depths (Dipre et al., 2018). In comparison, the subparallel stacking pattern of the stratified facies S4–S1 on top of the sediment waves (Figure 8g) indicates sedimentation in quieter environments less affected by bottom currents (Veeken & van Moerkerken, 2013).

5.2. Development of Glaciogenic Deposits

5.2.1. Trough Mouth Fan Contributions

Glaciogenic TMFs are typically formed in front of the cross-shelf glacial troughs—elongated, landward deepening bathymetric depressions on the broad continental margins (e.g., Batchelor & Dowdeswell, 2014; Ó Cofaigh et al., 2003). The relatively narrow Chukchi Rise extending offshore from the continental margin makes for a different geomorphic and depositional environment. A slope gradient is considered as one of the principal factors that control the development of TMFs (Batchelor & Dowdeswell, 2014; Ó Cofaigh et al., 2003). The upper slope gradient of the southwestern Chukchi Rise margin is less than 4° (2°–3° average) (Figure 1b), which is suggested to be close to a limit of an effective accumulation of glaciogenic material on the slope (e.g., Batchelor & Dowdeswell, 2014; Piper et al., 2012). Nevertheless, this gradient (~2°–3°) is considerably higher than the slope of well-developed TMFs (lower than 1°), such as on the Polar North Atlantic margins (e.g., Ó Cofaigh et al., 2003; Piper & Normark, 2009; Rydningen et al., 2015). The acoustically transparent sediment depocenters of subunits T4 and T3 on the southwestern middle to lower slope (Figures 3b, 3d, and 10a) indicate that the gravity-driven mass flow deposits (e.g., submarine landslides and debris flows) largely contributed to the TMF formation by filling the sediment accommodation space in the middle to lower slope, where the slope gradient is less than 1° (1,100–1,700 mwd: 0.9°; 1,700–2,200 mwd: 0.8°). This infilling may have led to a gradual decrease of the overall slope gradient (e.g., Faleide et al., 1996; O'Grady & Syvitski, 2002), which enabled the accumulation of glaciogenic debris lobes on the gentler slope of the southwestern Chukchi Rise margin. In contrast, a relatively high upper slope gradient (>4°) of the northwestern Chukchi Rise margin could be a reason for the poorly developed TMF (e.g., Batchelor & Dowdeswell, 2014; Ó Cofaigh et al., 2003; O'Grady & Syvitski, 2002; Piper & Normark, 2009).

Sediment supply for the TMF formation is also controlled by the subglacial geology of the continental shelf (Batchelor & Dowdeswell, 2014; Halberstadt et al., 2016; Ó Cofaigh et al., 2003, 2004; Solheim et al., 1998; Winsborrow et al., 2010). Prior deep seismostratigraphic data show that the preglacial strata with laterally continuous internal reflections are widely developed and completely cover high-standing crustal blocks of the Chukchi Rise (Dove et al., 2014; Hegewald & Jokat, 2013; Ilhan & Coakley, 2018) (Figure 1c). Weakly compacted marine sediment can be much easier eroded by grounded ice rather than crystalline rock or glaciogenic diamicton (e.g., Halberstadt et al., 2016; Moore, 1964; Ó Cofaigh et al., 2003; Wellner et al., 2001; Winsborrow et al., 2010). With respect to TMF development, the seaward-dipping bathymetric trough on the southwestern Chukchi Rise margin (Figure 1b) is relatively less developed than landward-dipping glacial troughs (Batchelor & Dowdeswell, 2014). A seaward-dipping trough can be easily eroded by repeated glacial advances despite a relatively short glacial history, as exemplified by the Mackenzie Trough at the Canadian Arctic margin (Batchelor et al., 2013). We infer that large quantities of sediment on the wide outer shelf of the southwestern Chukchi Rise margin were eroded by the fast-flowing grounded ice and remobilized downslope. This inference is consistent with the occurrence of large-scale glacial sediment T4/T3 depocenters on the midlower slope (Figures 4 and 10a).

Buried valleys reaching considerably larger dimensions are widely distributed across the Chukchi margin (Dove et al., 2014; Hill & Driscoll, 2008; Hill et al., 2007). Their occurrence on the Chukchi Rise at water depths of >300 m indicates their subglacial rather than river-born origin (Dove et al., 2014). The airgun seismic profiles at the western Chukchi Rise (e.g., Figure 5 in Dove et al., 2014) show buried incised channels likely formed by subglacial erosion. The eroded material may have contributed to the growth of the TMF. Conversely, the absence or poor development of TMFs at the northwestern shelf edge can be associated with a lack of large erosional features (Figure 7 in Dove et al., 2014).

Suspension settling from turbid meltwater plumes released from an ice-stream front can also contribute to the TMF growth and form gullies on the slope (Gales et al., 2013; Ó Cofaigh et al., 2003; Shipp et al., 1999). The development of gullies varies depending on the upper slope gradient (Klages et al., 2015; Ó Cofaigh et al., 2003). Gullies are weakly developed on the gentle (2° – 3°) southwestern slope of the Chukchi Rise, whereas deeply incised gullies occur on the steep ($>4^{\circ}$) northwestern slope (Figures 9a and 10b). This distribution implies that the TMF at the southwestern Chukchi Rise margin has been formed by complex sedimentary processes including debris flows and turbid meltwater plumes. Sedimentary deposits from turbid meltwater are characterized by acoustically stratified facies (Joe et al., 2020; Taylor et al., 2002), such as observed in our data between the debris flow packages (e.g., between T3b and T3a in Figure 4).

5.2.2. Glaciogenic Facies

Based on the mapped distribution of glaciogenic deposits, major contributions to the TMF at the southwestern Chukchi Rise margin can be attributed to the glacially fed debris flows of subunits T4/T3 and suspension settling from turbid meltwater plumes forming intercalated stratified facies (Figures 2–5). Contributions of T2 are much smaller, although the T2 grounding-zone wedge developed on the upper slope and outer shelf, while evidence from older glaciations has not been preserved (Figures 3f and 6). The distribution of glaciogenic deposits and an estimated position of the grounding line at the southwestern margin (Figure 10) indicate that the ice sheet at the time of T2 formation had the same or even larger extent than the previous T3/T4 glaciations. Therefore, changes in the glaciogenic inputs to the TMF likely had other controls. In general, the basal thermal condition of an ice sheet is a major control for sediment transport within a glacial system (Frederick et al., 2016; MacGregor et al., 2016; Menzies & Shilts, 2002). Larger quantities of sediment can be transported under temperate, wet than under cold polar/subpolar basal conditions (Alley et al., 1997; Benn & Evans, 2014; Davis et al., 2006; Kirkbride, 2002; Ottesen et al., 2005; Schomacker et al., 2010; Vorren & Laberg, 1997). In addition, proglacial sedimentation is strongly affected by the volume of meltwater and its proximity to the ice margin (C. H. Eyles & Eyles, 2010; Menzies & van der Meer, 2018). As a result, faster-flowing ice sheets with wet-based basal conditions could support more extensive sediment transportation and deposition at larger water depths. Another factor is that during the formation of T4/T3, large volumes of sediments were probably accumulated at the shelf break by subglacial transport of the readily erodible preglacial or ice-distal sediments, and subsequently remobilized downslope. In comparison, by the time of the T2 formation, most of the sediment on the outer shelf and upper slope may have been already consumed by the downslope transport. Furthermore, the older strata exposed on the shelf possibly became more compacted by ice-sheet loading (e.g., Böhm et al., 2009).

The continental slope of the western spur and the northwestern Chukchi Rise margin is mainly covered by well-developed stratified sediment layers (S4–S1) with insignificant lateral variation (Figures 3a, 3c, 3e, and 3g), except for the T1 accumulation at the northwestern outer shelf (Figures 3h and 7). This pattern indicates that gravity-driven mass flow processes were less active here than at the southwestern margin, probably due to colder basal conditions and/or lower sediment fluxes at a more northern location further away from the continental margin. The delivery of sediment to the western spur and the northwestern slope likely occurred mainly by iceberg-rafting and hemipelagic sedimentation, as common for proglacial glaciomarine environments (Ó Cofaigh et al., 2003). The stratified deposits of S4 (or S3 at some sites) to S1 in this area drape the preglacial slope features, such as sediment waves and gullies (Figures 8f, 8g, and 9a). This stratigraphy indicates that the active marine downslope and along-slope sedimentary processes were diminished since the onset of glaciations when depositional environments became predominantly glaciomarine.

The location of the major T1 glaciogenic depocenter on the outer shelf at relatively shallow water depths of <450 m (Figures 3h and 10a) suggests that T1 was deposited by a smaller and younger ice sheet than T2. Seismostratigraphic correlation of the boundary between U1 and U2 (Figure 2) further indicates that the formation of T1 postdated T2. This evidence is consistent with a lower thickness of acoustically stratified sediment on top of T1 at the northwestern margin (<1 m) than on T2 further south (>5 m) (Figures 3g, 6, and 7), resulting in well-expressed channels on top of T1 (Figures 7c and 10a). In comparison, the T2 surface features a hummocky proglacial moraine and well-developed gas seepage mounds indicative of a stronger glacial impact and/or longer postglacial period. We note, however, that the thickness of postglacial sediments depends not only on the time of deposition but also on sediment dynamics. It has been shown for

the study region that sedimentation rates since the last major glaciation overall decrease northwards (from >13 to <2 cm/kyr) indicating sediment sources further south (Schreck et al., 2018).

5.3. Past Ice-Sheet Dynamics and Related Processes

The fast-flowing ice streams marked in suitable seafloor sediments by the MSGL are primarily constrained by the ice sheet's mass balance, subglacial topography, and geology (e.g., Halberstadt et al., 2016). In general, the preserved MSGL indicates the latest grounded ice flow just before a final ice retreat, typically by floatation and breakup (e.g., Dowdeswell et al., 2008). The combined MBES and SBP data from the Chukchi Rise indicate that the deeper located (780–940 mwd), SSW–NNE oriented MSGL on the middle slope of the western spur were formed prior to the U2 deposits (Figures 8e and 9a). In comparison, N–S oriented MSGLs overriding T2 on the outer shelf were clearly formed by a younger and thinner ice mass, based on the overlying sediment thickness and the relationship between the water depth of the lineations and the required ice thickness (~114% of water depth) (Dowdeswell & Fugelli, 2012) (Figures 8d and 9a). This interpretation is consistent with the accumulation of acoustically stratified sediments with low backscatter intensity on top of the deeper MSGL, but only a thin sediment cover with higher backscatter intensity on top of the outer-shelf MSGL (Figures 8e, and 9a).

The general orientation of MSGL sets is subparallel to the direction of the continental shelf margin of the western Chukchi Rise (Figure 9a). This orientation may indicate that if the ice flow on the western Chukchi Rise was orthogonal to the shelf margin at the initial advance stage, it was later reoriented to slope-subparallel by a coalescence with the adjacent ice mass on the East Siberian margin. A similar, curved ice-flow pattern is known for the Ross Ice Shelf in the Antarctica, where several outlet glaciers along the coast of the Transantarctic Mountains (e.g., Byrd and Mulock Glaciers) merge and turn together into the main ice shelf (LeDoux et al., 2017; Rignot et al., 2011). Regardless of whether the MSGLs were formed by a coherent ice stream or an armada of large icebergs derived from a collapsing ice sheet, it can be assumed that the overall ice flow at the Chukchi–East Siberian margin was directed toward the central Arctic Ocean.

The sets of laterally extended, isobath-parallel recessional moraines on the outer shelf to upper slope of the western margin are apparently associated mostly with the T2 deposits (Figures 8d and 9a). A prominent stand-alone ridge at shallower water depths of ~400–420 m might indicate a younger glacial event (Figure 9a). This ridge overrides the MSGL in this area and thus was formed after the ice-streaming event marked by the MSGL. Overall the ridges likely indicate a stepwise upslope retreat of the ice-sheet grounding line (Batchelor et al., 2018; Jakobsson et al., 2008; Ottesen & Dowdeswell, 2006; Polyak et al., 2001) controlled by the rising sea levels during deglaciation(s) (Cronin et al., 2017). This pattern contrasts with the bathymetry independent MSGL controlled by the ice-sheet mass balance and topography during ice-sheet collapsing events. The ice-sheet collapses and retreats on the Chukchi Rise may have been triggered by the instability of an ice shelf (shelves) over the adjacent basins in the absence of pinning points (e.g., Halberstadt et al., 2016; Jakobsson et al., 2016; Wellner et al., 2001). Different patterns of the recessional ridge fields on the western spur and the southwestern margin may reflect a lateral variability in the retreat rapidity (Figures 9a and 10b). The western spur features multiple recessional ridges with a high lateral continuity, indicating a relatively gradual, stepwise retreat (e.g., Dowdeswell et al., 2008) (Figures 9a and 10b). The sparser, less developed recessional moraines on the outer-shelf terrace of the southwestern margin possibly indicate a more abrupt retreat of the grounding line from the shelf edge to the inner shelf (Figures 6, 9a, and 10b). The faster retreat may have been facilitated by a deeper shelf break and a flatter gradient of the shelf-edge terrace in the southwestern area (Figures 9a and 10b).

The area of hummocky moraines and mounds developed on the southwestern outer shelf and upper slope is covered by a moderately thick acoustically stratified sediment of S1 with relatively low backscatter intensity (Figures 8b and 9a). This pattern indicates a farther/earlier shelfward retreat of the grounded ice sheet in this area than further north on the western spur and the northwestern margin featuring thinner overlying stratified sediments and higher backscatter intensity (Figures 9a and 10b). The seepage-related mound structures preserved on the outer shelf and upper slope of the southwestern margin also pre-date the U1 deposits (Kim et al., 2020) (Figure 8b), thus indicating no ice-sheet readvance in this area after the T2 formation (Figure 10b).

Based on the stratigraphic position of the buried incised surface and the thickness of the overlying acoustically stratified sediments, the gullies on the northwestern slope (Figures 8f and 9a) are probably older (pre-U3) than at the southwestern margin (U2; Figures 8a and 9a). The gully formation can be explained by a grounded ice advance to the shelf edge that provided turbid meltwater pulses capable of incising canyons on the steep slope (A. L. Lowe & Anderson, 2003; Ó Cofaigh et al., 2003). After the gully formation event at the northwestern slope, no sufficient sediment-laden meltwater was generated in this area, possibly due to a change in the basal thermal condition from the temperate-wet to the polar-cold (Rebesco & Camerlenghi, 2008). The younger and smaller gullies on the southwestern slope indicate overall lower amounts of subglacial meltwater discharged during the T2 glacial event. In comparison to the deeper located slope gullies, the fresh-looking, relatively shallow but broad (up to 1,000 m) channels on top of the T1 wedge on the northwestern shelf likely have a different process of formation. Considering their geometry and the position on a gently dipping surface of the youngest glacial deposit, they may be related to the subglacial drainage at the final deglacial stages. On the other hand, low backscatter intensities of the T1 wedge surface (Figure 7c) are more common for proglacial rather than subglacial deposits. Clarifying the nature of this glaciogenic accumulation requires denser data coverage, including shelf-to-slope lines.

The preglacial sediment waves expressed in bathymetry-parallel, undulating morphology are well preserved on the slope of the western spur and the northwestern margin due to the conformable geometry of the overlying acoustically stratified deposits of S4–S1 (Figure 8g). The absence of similar undulations on the southwestern slope (Figure 9a) can be explained by the burial of preglacial sediment waves by the thick TMF deposits (Dove et al., 2014; Hegewald & Jokat, 2013; Ilhan & Coakley, 2018).

5.4. Glaciation History

5.4.1. Ice-Sheet Distribution and Provenance

The overall distribution of glaciogenic seafloor morphology and sediment stratigraphy in the study area is consistent with the notion that Arctic ice sheets advanced from the continental margins toward the central Arctic Ocean and retreated upslope back to the margins and shallow bathymetric areas (Dove et al., 2014; Engels et al., 2008; Jakobsson et al., 2008, 2010, 2014; Polyak et al., 2001, 2007). If the entire Arctic Ocean was covered by a thick ice shelf during some of the peak glaciations, it likely behaved as a single large ice mass outflowing into the North Atlantic (Hughes et al., 1977; Jakobsson et al., 2016). Due to these changes in ice flow and a lack of major topographic constraints, such as cross-shelf troughs or interisland channels, the directions of ice advances and retreats at the western Chukchi Rise margin are more complex than on the well-developed glaciated margins around the Antarctic, Polar North Atlantic, and the Canadian Arctic Archipelago (Batchelor & Dowdeswell, 2014; Batchelor et al., 2014; Halberstadt et al., 2016; Ó Cofaigh et al., 2003; Winsborrow et al., 2010).

Prior MBES data from the Chukchi Borderland collected mostly east of the study area indicate a prevalent SE–NW to ESE–WNW trending impact of grounded ice mass(es) projected to originate from the northwestern sector of the Laurentide Ice Sheet (Figure 1a) (Dove et al., 2014; Engels et al., 2008; Jakobsson et al., 2008, 2014; Polyak et al., 2001, 2007). Another distinct set of directional seafloor features on the eastern side of the Chukchi Rise appears to irradiate from the Chukchi Shelf (Dove et al., 2014; Jakobsson et al., 2014), as also observed in our ancillary data (Figure 9c). The cooccurrence of these major flowline trends makes a complex picture reflecting the interaction of different ice flows. The evidence for ice provenance in our seismostratigraphic and geomorphic data from the western side of the Chukchi Rise provides another important key to this complex glacial history.

Based on the TMF location in the seaward-dipping bathymetric trough at the southwestern Chukchi Rise margin and the distribution of glaciogenic deposits T4/T3 preserved on the slope, the general ice-flow direction in this area was probably northwestward from the Chukchi Shelf toward the Chukchi Basin (Figure 10a). A similar, more S–N orientation of a MSGSL set on the outer shelf formed apparently on top of T2 (Figure 9a) also suggests an ice flow from the Chukchi margin. The prominent grounding-zone wedges on the western margin at the shelf edge and further shelfward probably mark standstill positions of the grounded ice margins (Figure 10b). The upslope deglacial retreat of the ice margins is marked by sets of recessional moraines (Figures 8d, 9a, and 10b), also observed at the northern Chukchi Rise tip (Figure 9b).

and in prior data from the Chukchi Rise and Cap further east and north (Dove et al., 2014; Jakobsson et al., 2008). In our data, these ridges appear to be primarily associated with T2. Similar grounding-zone standstill and retreat features for T4/T3 apparently have not been preserved.

The overall picture emerging from these data indicates that ice coming from the Laurentide Ice Sheet may have been deflected northwards (i.e., toward the Canada Basin) by the ice sheet(s) irradiating from the Chukchi margin. Alternatively, it can also be inferred that the younger Chukchi-centered ice sheets overprinted the depositional/geomorphic evidence of the older ice masses with different flow trajectories. Considering that our data cover four grounding events with a long glacial history since the mid-Pleistocene, it is more likely that coeval ice masses with different provenance actually interacted on the Chukchi Rise. We note that while multiple seafloor features indicate ice flow(s) from the Chukchi continental margin, they cannot be confirmed by data from the shelf itself (Dove et al., 2014; Jakobsson et al., 2014). The obvious reason for this lack of evidence is that sedimentary bedforms cannot preserve on the shallow, current-swept Chukchi Shelf. Deeper geophysical records and drilling boreholes are needed to resolve this gaping blind spot.

On the western side of the Chukchi Rise, the ice-flow interactions can also be envisaged with the marine-based ice sheet on the Chukchi–East Siberian margin (Dove et al., 2014; Niessen et al., 2013). While still sketchily understood (Batchelor et al., 2019, and references therein), this ice sheet is corroborated by multiple seafloor data (Dove et al., 2014; Jakobsson et al., 2016; Joe et al., 2020; Niessen et al., 2013; O'Regan et al., 2017; Schreck et al., 2018) and paleoclimatic modeling studies (Colleoni et al., 2016; Gasson et al., 2018). If the ice shelves extending from the East Siberian and Chukchi margins consequently coalesced and started to behave like a single large ice mass (Hughes et al., 1977; Jakobsson et al., 2016), the flow direction would have been reoriented toward the central Arctic Ocean (yellow arrows in Figure 10b), orthogonally to the Chukchi–East Siberian margin. The overall ice flow on the western Chukchi Rise may have accounted for the SSW–NNE trending, deep-sited MSGL on top of the T3 deposits on the western spur (Figure 10b). The large water depth of these features (>900 m) is similar to the ice grounding on the Lomonosov Ridge in the center of the Arctic Ocean (Jakobsson et al., 2001, 2008, 2010, 2014; Polyak et al., 2001). Elaborating on these mechanisms is important for understanding the behavior of the Arctic glaciations and, more generally, the large marine-based ice complexes.

5.4.2. Age Framework

While there are no direct constraints for the age of the mapped glaciogenic deposits, a tentative age framework can be outlined from stratigraphic data available from the adjacent areas (Dipre et al., 2018; Joe et al., 2020; Polyak et al., 2007; Schreck et al., 2018; Wang et al., 2013). In particular, combined geological/geophysical data from the southeastern part of the Chukchi Borderland (ramp to the Northwind Ridge; Figure 1b) provide age constraints for the last two glacial events impacting this seafloor area (Polyak et al., 2007). Glacial diamicton dated to the Last Glacial Maximum (LGM) of the Marine Isotope Stage (MIS) 2, ca. 15–25 ka, is associated with the W–E trending MSGL at modern water depths to ~420–430 m (Polyak et al., 2007). A nearby MSGL field mapped at the southeastern side of the Chukchi Rise and presented in this study has similar bathymetric and geomorphic characteristics (Figures 1b and 9c). The MSGL orientations in these two fields indicate ice streaming from the Chukchi margin. This picture is consistent with the distribution of the T1 deposits at the western Chukchi Rise, which is restricted to water depths of <450 m and are covered with only thin postglacial sediments (Figures 3h, 7, and 10a). Well-expressed channels on top of the T1 deposit at the northwestern shelf possibly represent the proglacial or subglacial drainage during deglaciation, similar to, but younger than buried valleys reported for the Chukchi Rise by Dove et al. (2014). These characteristics of the T1 deposits allow for attribution of their formation to the LGM. This interpretation is also consistent with sediment-core evidence from the Chukchi Basin that suggests the presence of the LGM grounded ice masses at shallower depths in this region (Joe et al., 2020; Schreck et al., 2018; Wang et al., 2013).

Glacial deposits attributed to MIS 4 (ca. 60–70 ka) appear to be widely distributed in the study region, including tills on the bathymetric highs, debris lobes on the slopes, and thick glaciomarine deposits in the basins (Joe et al., 2020; Polyak et al., 2007; Schreck et al., 2018; Wang et al., 2013). The corresponding glaciogenic bedforms such as MSGL and morainic ridges extend to the shelf edge at ~600–700 mwd (Dove et al., 2014; Jakobsson et al., 2008; Polyak et al., 2007). At the southern Northwind Ridge, these deposits feature the

SE–NW trending MSGL indicative of an ice flow from the Laurentide Ice Sheet (Polyak et al., 2007). MSGLs with a similar orientation and depth range are also widely reported elsewhere from the Northwind Ridge, Chukchi Cap, and Alaskan margin (Dove et al., 2014; Engels et al., 2008; Jakobsson et al., 2005, 2008). The seismostratigraphic and bathymetric position of the T2 deposits indicates their probable relation to the coeval glacial event. The seismostratigraphic boundaries of U2, including T2 (H2/H3, Figure 2a), apparently correspond to reflectors R2/R3 defined for the East Siberian (Arlliss Plateau) slope across the Chukchi Basin (Joe et al., 2020). These reflectors were related to prominent IRD layers formed by the iceberg discharge pulses from the Laurentide Ice Sheet during the deglaciations following and preceding MIS 4. Despite probable synchronicity with a Laurentide-sourced ice advance inferred for more eastern areas, this glaciation on the western Chukchi Rise apparently had a different provenance based on the S–N-trending orientation of the MSGL on top of T2. According to this orientation, the main grounded ice was located south of the Chukchi Rise at least at the late glacial stages. This inference is consistent with the formation of a hummocky moraine south of the MSGL field (Figure 9a).

The age controls for the older glacial deposits T4/T3 preserved in the study area only as debris lobes on the slope are more speculative. The potentially largest glacial impact in the Arctic has been inferred for MIS 6 (late Middle Pleistocene, ca. 130–190 ka) based on a very deep (>1 km) ice grounding on the Lomonosov Ridge in the center of the Arctic Ocean constrained to this glacial interval (Jakobsson et al., 2001, 2010, 2014, 2016; Polyak et al., 2001). Reconstructed circum-Arctic ice-sheet limits also show a very large extent for the northern North America and the largest for the northern Eurasia (Batchelor et al., 2019, and references therein). Glacial diamicton probably of this age has been recovered on top of the northern part of the Northwind Ridge in front of a prominent morainic ridge (Dipre et al., 2018; Jakobsson et al., 2010). A large pre-MIS 4 debris lobe on the East Siberian slope (Arlliss Plateau: Joe et al., 2020) may also belong to the MIS 6. A possible attribution of the glaciogenic T3 deposits to this glaciation is indirectly supported by the MSGL set formed at water depths to >900 m on top of T3 and aligned with the direction toward the central Arctic Ocean (Figures 9a and 10). Two glaciogenic lobes identified as subunits of T3 (Figure 4) indicate two glacial events, possibly stages of the same glaciation.

The age of the oldest glaciogenic unit T4 can be considered in relation to the history of a large-scale glacial erosion on the Chukchi margin/borderland. Based on regional airgun seismic data, the glaciogenic erosional boundary occurs on top of deposits overlying the late Miocene unconformity and is broadly attributed to Plio-Pleistocene (Hegewald, 2012; Hegewald & Jokat, 2013; İlhan & Coakley, 2018). In sediment cores from the northern Northwind Ridge the contact between preglacial and glaciogenic strata is constrained by cyclostratigraphy and strontium isotope dating to MIS 16/20, ca. 0.7–0.9 Ma (Dipre et al., 2018; Polyak et al., 2013). This timing corresponds to a prominent change in paleoclimatic conditions (mid-Pleistocene Transition) related to a shift in the prevailing orbital cyclicities and the development of huge ice sheets in the Northern Hemisphere in the Middle Pleistocene starting with MIS 16 (Clark et al., 2006; Lisiecki & Raymo, 2005).

Based on the airgun seismic data, there is one glaciogenic sedimentary package between the reflector H5 (lower boundary of unit U4) and the initial glaciogenic unconformity (Dove et al., 2014) (H6; Figure S3). This stratigraphy indicates that T4 was formed during the second regional glacial event. If the first glacial expansion occurred in MIS 16 or somewhat earlier, the T4 age can be placed somewhere in the lower part of the Middle Pleistocene, such as MIS 12 that was a prominent glaciation based on a sedimentary record from the Canada Basin (Dong et al., 2017).

Constraining the outlined stratigraphic estimates more conclusively requires more sediment cores, preferably with long records such as pursued by the International Ocean Discovery Program (IODP). Our data may contribute useful information for developing an IODP project in the Chukchi region.

6. Summary and Conclusions

This study presents the detailed, high-resolution SBP and MBES data from the Chukchi Rise in the western Arctic Ocean. The combined seismostratigraphic and morphobathymetric analysis sheds new light on the past ice-sheet/ice-shelf distribution and dynamics and related sedimentary/geomorphic processes. Results

of this study provide important empirical constraints for reconstructing Quaternary glaciations in and around the Arctic Ocean.

Based on the acoustic character of sediments revealed by the SBP data, we identify four seismostratigraphic units, U4–U1, each including acoustically stratified and transparent facies classified as subunits S4–S1 and T4–T1, respectively. The transparent facies are interpreted as glaciogenic sediments deposited during glacial events probably spanning most of the Middle to Late Pleistocene (ca. 0.5–1 Ma). The older transparent deposits (T4/T3) are preserved on the southwestern midlower slope of the Chukchi Rise as debris lobes contributing to a large TMF identified on the regional airgun seismic records. On the upper slope to outer shelf, these features are replaced by a younger, wedge- to sheet-shaped deposit (T2) with an erosional lower boundary and glaciogenic bedforms at the surface, such as MSGLs, recessional morainic ridges, and hummocky moraines. The main T2 deposit at the shelf break is interpreted as a glacial grounding-zone wedge possibly formed during MIS 4 (ca. 60–70 ka). The youngest transparent deposit (T1), attributed to the MIS 2 (ca. 15–25 ka), occurs on the outer shelf at shallower water depths.

Two mapped sets of MSGL trending SSW–NNE on top of T3 on the middle slope and S–N on T2 on the outer shelf demonstrate at least two fast ice-streaming events of different ages. The older, deeper sited (ca 780–940 mwd) MSGL may be related to a passage of a large ice shelf formed over the western Arctic Ocean. Contour-parallel, nested recessional-moraine ridges are identified along the outer shelf to upper slope on top of the grounding-zone deposits. Correlative ridges are also mapped north of the main study area, thus indicating their distribution along the entire Chukchi Rise margin. The well-developed ridges mark a step-wise retreat of the grounded ice margin, likely controlled by rising sea levels during deglaciation(s). More random ridges at a flatter bathymetric bench on the southwestern margin may indicate a faster retreat. The different orientations of ice advances and retreats reflect a complex geomorphic setting of the borderland that lacks major topographic controls typical for glaciated continental margins, such as cross-shelf glacial troughs.

The seismostratigraphic and geomorphic data suggest that ice-flow directions for all of the identified glacial events were principally from the Chukchi continental margin south of the study area. Based on the additional mapping and prior data, the same ice-sheet origin can be projected for glaciogenic bedforms on the eastern side of the Chukchi Rise, along with the evidence of ice flow(s) apparently originating from the northwestern Laurentide Ice Sheet. This complex picture shows that the Chukchi Rise was an area of intense interaction(s) of different ice sheets/ice shelves that affected the overall glaciation development in the Arctic Ocean.

In addition to glaciogenic deposits and bedforms, our data identify a number of related or independent seabed features including mounds, gullies/channels, and sediment waves. Mounds, presumably formed by gas seepage, grow from the surface of the T2 deposit, possibly triggered by glaciogenic pressure effects. Buried gullies on the upper slope were likely incised by proglacial turbid water flows, while fresh-looking, broad channels on top of the T1 deposit on the shelf may be related to the proglacial or subglacial drainage during the last deglaciation. Sediment waves are characteristic for preglacial sediments on the middle slope, thus indicating active hydrodynamic conditions at the intermediate water depths before the onset of major glaciations.

Acknowledgments

The authors are grateful to the captain and crew of the IBRV *Araon* for their help provided during the Arctic expeditions since 2012. This work was supported by the Korea Polar Research Institute (KOPRI) grant PM20050 (KIMST grant 20160247). Y. J. Joe and S.-I. Nam acknowledge funding from the KOPRI grant PE20350. F. Niessen was funded by the AWI Research Program PACES-II, Workpackages 3.1 and 3.2. We thank SeisWare Inc. for the academic grant program for using the software for seismostratigraphic analysis and mapping. The authors thank two anonymous reviewers for providing many helpful comments and suggestions for improving the manuscript.

Data Availability Statement

SBP data processing, MBES data gridding, and mapping were conducted using the open-source software Seismic Unix (<https://wiki.seismic-unix.org>), Generic Mapping Tools (<https://www.generic-mapping-tools.org>), and QGIS (<https://qgis.org>). The MBES bathymetry/backscatter and the seismostratigraphic units/subunits thickness mapping results used in this study are archived by the Korea Polar Data Center and are available at <https://kpdc.kopri.re.kr>.

References

- Alley, R. B., Anandakrishnan, S., Dupont, T. K., Parizek, B. R., & Pollard, D. (2007). Effect of sedimentation on ice-sheet grounding-line stability. *Science*, 315(5820), 1838–1841.

- Alley, R. B., Blankenship, D. D., Bentley, C. R., & Rooney, S. T. (1986). Deformation of till beneath ice stream B, West Antarctica. *Nature*, 322(6074), 57–59.
- Alley, R. B., Cuffey, K. M., Evenson, E. B., Strasser, J. C., Lawson, D. E., & Larson, G. J. (1997). How glaciers entrain and transport basal sediment: Physical constraints. *Quaternary Science Reviews*, 16(9), 1017–1038. [https://doi.org/10.1016/S0277-3791\(97\)00034-6](https://doi.org/10.1016/S0277-3791(97)00034-6)
- Anderson, J. B. (1999). Antarctic marine geology (Vol. 289). Cambridge: Cambridge University Press.
- Batchelor, C. L., & Dowdeswell, J. A. (2014). The physiography of High Arctic cross-shelf troughs. *Quaternary Science Reviews*, 92, 68–96. <https://doi.org/10.1016/j.quascirev.2013.05.025>
- Batchelor, C. L., & Dowdeswell, J. A. (2015). Ice-sheet grounding-zone wedges (GZWs) on high-latitude continental margins. *Marine Geology*, 363, 65–92. <https://doi.org/10.1016/j.margeo.2015.02.001>
- Batchelor, C. L., Dowdeswell, J. A., & Ottesen, D. (2018). Submarine glacial landforms. In A. Micallef, S. Krastel, & A. Savini (Eds.), *Submarine geomorphology* (pp. 207–234). Cham, Switzerland: Springer International Publishing. https://doi.org/10.1007/978-3-319-57852-1_12
- Batchelor, C. L., Dowdeswell, J. A., & Pietras, J. T. (2013). Seismic stratigraphy, sedimentary architecture and palaeo-glaciology of the MacKenzie Trough: Evidence for two Quaternary ice advances and limited fan development on the western Canadian Beaufort Sea margin. *Quaternary Science Reviews*, 65, 73–87. <https://doi.org/10.1016/j.quascirev.2013.01.021>
- Batchelor, C. L., Dowdeswell, J. A., & Pietras, J. T. (2014). Evidence for multiple Quaternary ice advances and fan development from the Amundsen Gulf cross-shelf trough and slope, Canadian Beaufort Sea margin. *Marine and Petroleum Geology*, 52, 125–143. <https://doi.org/10.1016/j.marpetgeo.2013.11.005>
- Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., et al. (2019). The configuration of Northern Hemisphere ice sheets through the Quaternary. *Nature Communications*, 10(1), 3713. <https://doi.org/10.1038/s41467-019-11601-2>
- Batchelor, C. L., Ottesen, D., & Dowdeswell, J. A. (2017). Quaternary evolution of the northern North Sea margin through glacially derived flow and contourite deposition. *Journal of Quaternary Science*, 32(3), 416–426. <https://doi.org/10.1002/jqs.2934>
- Benn, D., & Evans, D. J. (2014). *Glaciers and glaciation* (2nd ed.). Hodder Education.
- Bennett, M. R., Huddart, D., Hambrey, M. J., & Ghiene, J. F. (1996). Moraine development at the High-Arctic valley glacier Pedersenbreen, Svalbard. *Geografiska Annaler - Series A: Physical Geography*, 78(4), 209–222. <https://doi.org/10.1080/04353676.1996.11880468>
- Böhm, G., Ocañoğlu, N., Picotti, S., & De Santis, L. (2009). West Antarctic Ice Sheet evolution: New insights from a seismic tomographic 3D depth model in the Eastern Ross Sea (Antarctica). *Marine Geology*, 266(1), 109–128.
- Boone, S. J., & Eyles, N. (2001). Geotechnical model for great plains hummocky moraine formed by till deformation below stagnant ice. *Geomorphology*, 38(1), 109–124. [https://doi.org/10.1016/S0169-555X\(00\)00072-6](https://doi.org/10.1016/S0169-555X(00)00072-6)
- Bradwell, T., Stoker, M. S., Gollidge, N. R., Wilson, C. K., Merritt, J. W., Long, D., et al. (2008). The northern sector of the last British Ice Sheet: Maximum extent and demise. *Earth-Science Reviews*, 88(3), 207–226. <https://doi.org/10.1016/j.earscirev.2008.01.008>
- Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., et al. (2006). The middle Pleistocene transition: Characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂. *Quaternary Science Reviews*, 25(23), 3150–3184. <https://doi.org/10.1016/j.quascirev.2006.07.008>
- Coakley, B. J., Ilhan, I., & Chukchi Edges Science Party. (2011). Chukchi Edges Project—Geophysical constraints on the history of the Amerasia Basin. In *AGU Fall Meeting Abstracts* (Vol. 1, p. 2365).
- Colleoni, F., Kirchner, N., Niessen, F., Quiquet, A., & Liakka, J. (2016). An East Siberian ice shelf during the Late Pleistocene glaciations: Numerical reconstructions. *Quaternary Science Reviews*, 147, 148–163.
- Cronin, T. M., O'Regan, M., Pearce, C., Gemery, L., Toomey, M., Semiletov, I., & Jakobsson, M. (2017). Deglacial sea level history of the East Siberian Sea and Chukchi Sea margins. *Climate of the Past*, 13(9), 1097–1110. <https://doi.org/10.5194/cp-13-1097-2017>
- Darby, D. A., Jakobsson, M., & Polyak, L. (2005). Icebreaker expedition collects key Arctic seafloor and ice data. *Eos, Transactions American Geophysical Union*, 86(52), 549–552.
- Davis, P. T., Briner, J. P., Coulthard, R. D., Finkel, R. W., & Miller, G. H. (2006). Preservation of Arctic landscapes overridden by cold-based ice sheets. *Quaternary Research*, 65(1), 156–163. <https://doi.org/10.1016/j.yqres.2005.08.019>
- DeConto, R. M., & Pollard, D. (2016). Contribution of Antarctica to past and future sea-level rise. *Nature*, 531(7596), 591–597.
- Dipre, G. R., Polyak, L., Kuznetsov, A. B., Oti, E. A., Ortiz, J. D., Brachfeld, S. A., et al. (2018). Plio-Pleistocene sedimentary record from the Northwind Ridge: New insights into paleoclimatic evolution of the western Arctic Ocean for the last 5 Ma. *Arktos*, (1), 24.
- Donda, F., O'Brien, P. E., De Santis, L., Rebesco, M., & Brancolini, G. (2008). Mass wasting processes in the Western Wilkes Land margin: Possible implications for East Antarctic glacial history. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 260(1), 77–91.
- Dong, L., Liu, Y., Shi, X., Polyak, L., Huang, Y., Fang, X., et al. (2017). Sedimentary record from the Canada Basin, Arctic Ocean: Implications for late to middle Pleistocene glacial history. *Climate of the Past*, 13(5), 511–531. <https://doi.org/10.5194/cp-13-511-2017>
- Dove, D., Polyak, L., & Coakley, B. (2014). Widespread, multi-source glacial erosion on the Chukchi margin, Arctic Ocean. *Quaternary Science Reviews*, 92, 112–122.
- Dowdeswell, J. A., & Bamber, J. L. (2007). Keel depths of modern Antarctic icebergs and implications for sea-floor scouring in the geological record. *Marine Geology*, 243(1), 120–131.
- Dowdeswell, J. A., Canals, M., Jakobsson, M., Todd, B. J., Dowdeswell, E. K., & Hogan, K. A. (2016). In J. A. Dowdeswell, M. Canals, M. Jakobsson, B. J. Todd, E. K. Dowdeswell, & K. A. Hogan (Eds.), *Atlas of submarine glacial landforms: Modern, Quaternary and ancient* (Vol. 46). London: Geological Society of London.
- Dowdeswell, J. A., Cofaigh, C. Ó., & Pudsey, C. J. (2004). Thickness and extent of the subglacial till layer beneath an Antarctic paleo-ice stream. *Geology*, 32(1), 13–16.
- Dowdeswell, J. A., & Fugelli, E. M. G. (2012). The seismic architecture and geometry of grounding-zone wedges formed at the marine margins of past ice sheets. *The Geological Society of America Bulletin*, 124(11–12), 1750–1761. <https://doi.org/10.1130/B30628.1>
- Dowdeswell, J. A., Kenyon, N. H., Elverhøi, A., Laberg, J. S., Hollender, F.-J., Mienert, J., & Siegert, M. J. (1996). Large-scale sedimentation on the glacier-influenced Polar North Atlantic margins: Long-range side-scan sonar evidence. *Geophysical Research Letters*, 23(24), 3535–3538.
- Dowdeswell, J. A., Ottesen, D., Evans, J., Cofaigh, C. Ó., & Anderson, J. B. (2008). Submarine glacial landforms and rates of ice-stream collapse. *Geology*, 36(10), 819–822.
- Dowdeswell, J. A., & Siegert, M. J. (1999). Ice-sheet numerical modeling and marine geophysical measurements of glacier-derived sedimentation on the Eurasian Arctic continental margins. *The Geological Society of America Bulletin*, 111(7), 1080–1097.
- Elvenes, S., & Dowdeswell, J. A. (2016). Possible 'lift-off moraines' at grounded ice-sheet margins, North Norwegian shelf edge. *Geological Society, London, Memoirs*, 46(1), 247–248.
- Elverhøi, A., Dowdeswell, J. A., Funder, S., Mangerud, J., & Stein, R. (1998). Glacial and oceanic history of the Polar North Atlantic margins: An overview. *Quaternary Science Reviews*, 17(1), 1–10. [https://doi.org/10.1016/S0277-3791\(97\)00073-5](https://doi.org/10.1016/S0277-3791(97)00073-5)

- Engels, J. L., Edwards, M. H., Polyak, L., & Johnson, P. D. (2008). Seafloor evidence for ice shelf flow across the Alaska–Beaufort margin of the Arctic Ocean. *Earth Surface Processes and Landforms*, 33(7), 1047–1063. <https://doi.org/10.1002/esp.1601>
- Eyles, C. H., & Eyles, N. (2010). Glacial deposits. In *Facies models* (Vol. 4, pp. 73–104). Newfoundland, Canada: The Geological Association of Canada.
- Eyles, C. H., Eyles, N., & Lagoe, M. B. (1991). The Yakataga Formation: A six million year record of temperate glacial marine sedimentation in the Gulf of Alaska. In J. B. Anderson & G. M. Ashley (Eds.), *Glacial marine sedimentation: Paleoclimatic significance* (Vol. 261, pp. 159–180). Boulder, CO: Geological Society of America.
- Eyles, N., Boyce, J. I., & Barendregt, R. W. (1999). Hummocky moraine: Sedimentary record of stagnant Laurentide Ice Sheet lobes resting on soft beds. *Sedimentary Geology*, 123(3), 163–174. [https://doi.org/10.1016/S0037-0738\(98\)00129-8](https://doi.org/10.1016/S0037-0738(98)00129-8)
- Faleide, J. I., Solheim, A., Fiedler, A., Hjelstuen, B. O., Andersen, E. S., & Vanneste, K. (1996). Late Cenozoic evolution of the western Barents Sea–Svalbard continental margin. *Global and Planetary Change*, 12(1), 53–74.
- Fransner, O., Noormets, R., Flink, A. E., Hogan, K. A., & Dowdeswell, J. A. (2018). Sedimentary processes on the continental slope off Vitøya and Albertini troughs north of Nordaustlandet, Svalbard—The importance of structural–geological setting in trough-mouth fan development. *Marine Geology*, 402, 194–208. <https://doi.org/10.1016/j.margeo.2017.10.008>
- Frederick, B. C., Young, D. A., Blankenship, D. D., Richter, T. G., Kempf, S. D., Ferraccioli, F., & Siegert, M. J. (2016). Distribution of subglacial sediments across the Wilkes Subglacial Basin, East Antarctica. *Journal of Geophysical Research: Earth Surface*, 121, 790–813. <https://doi.org/10.1002/2015JF003760>
- Gales, J. A. (2013). *The geomorphology of Antarctic submarine slopes* (PhD thesis). Manchester, UK: The University of Manchester.
- Gales, J. A., Larter, R. D., Leat, P. T., & Jokat, W. (2016). Components of an Antarctic trough-mouth fan: Examples from the Crary Fan, Weddell Sea. In J. A. Dowdeswell, M. Canals, M. Jakobsson, B. J. Todd, E. K. Dowdeswell, & K. A. Hogan (Eds.), *Atlas of submarine glacial landforms: Modern, Quaternary and ancient* (Vol. 46, pp. 377–378). London: Geological Society, London, Memoirs.
- Gales, J. A., Larter, R. D., Mitchell, N. C., & Dowdeswell, J. A. (2013). Geomorphic signature of Antarctic submarine gullies: Implications for continental slope processes. *Marine Geology*, 337, 112–124.
- Gasson, E. G. W., DeConto, R. M., Pollard, D., & Clark, C. D. (2018). Numerical simulations of a kilometre-thick Arctic ice shelf consistent with ice grounding observations. *Nature Communications*(1), 1510.
- Gebhardt, A. C., Jokat, W., Niessen, F., Matthiessen, J., Geissler, W. H., & Schenke, H. W. (2011). Ice sheet grounding and iceberg plow marks on the northern and central Yermak Plateau revealed by geophysical data. *Quaternary Science Reviews*, 30(13), 1726–1738. <https://doi.org/10.1016/j.quascirev.2011.03.016>
- Gulick, S. P. S., Shevenell, A. E., Montelli, A., Fernandez, R., Smith, C., Warny, S., et al. (2017). Initiation and long-term instability of the East Antarctic Ice Sheet. *Nature*, 552, 225–229.
- Halberstadt, A. R. W., Simkins, L. M., Greenwood, S. L., & Anderson, J. B. (2016). Past ice-sheet behaviour: Retreat scenarios and changing controls in the Ross Sea, Antarctica. *The Cryosphere*, 10(3), 1003–1020. <https://doi.org/10.5194/tc-10-1003-2016>
- Hall, J. K. (1990). Chukchi Borderland. In A. Grantz, L. Johnson, & J. F. Sweeney (Eds.), *The Arctic Ocean region* (Vol. L, pp. 337–350). Boulder, CO: Geological Society of America.
- Hegewald, A. (2012). *The Chukchi region–Arctic Ocean–Tectonic and sedimentary evolution* (PhD thesis). Jena, Germany: Chemistry and Geoscience University of Jena.
- Hegewald, A., & Jokat, W. (2013). Tectonic and sedimentary structures in the northern Chukchi region, Arctic Ocean. *Journal of Geophysical Research: Solid Earth*, 118, 3285–3296. <https://doi.org/10.1002/jgrb.50282>
- Hill, J. C., & Driscoll, N. W. (2008). Paleodrainage on the Chukchi shelf reveals sea level history and meltwater discharge. *Marine Geology*, 254(3), 129–151. <https://doi.org/10.1016/j.margeo.2008.05.018>
- Hill, J. C., Driscoll, N. W., Brigham-Grette, J., Donnelly, J. P., Gayes, P. T., & Keigwin, L. (2007). New evidence for high discharge to the Chukchi shelf since the Last Glacial Maximum. *Quaternary Research*, 68(2), 271–279.
- Himmeler, T., Sahy, D., Martma, T., Bohrmann, G., Plaza-Faverola, A., Bünz, S., et al. (2019). A 160,000-year-old history of tectonically controlled methane seepage in the Arctic. *Science Advances*, 5(8), eaaw1450. <https://doi.org/10.1126/sciadv.aaw1450>
- Howat, I. M., Joughin, I., & Scambos, T. A. (2007). Rapid changes in ice discharge from Greenland outlet glaciers. *Science*, 315(5818), 1559–1561. <https://doi.org/10.1126/science.1138478>
- Huang, Z., Siwabessy, J., Cheng, H., & Nichol, S. (2018). Using multibeam backscatter data to investigate sediment-acoustic relationships. *Journal of Geophysical Research: Oceans*, 123, 4649–4665. <https://doi.org/10.1029/2017JC013638>
- Hughes, T., Denton, G., & Grosswald, M. (1977). Was there a late-Würm Arctic ice sheet? *Nature*, 266(5603), 5967602.
- Hunkins, K., Herron, T., Kutschale, H., & Peter, G. (1962). Geophysical studies of the Chukchi Cap, Arctic Ocean. *Journal of Geophysical Research*, 67(1), 235–247.
- Ilhan, I., & Coakley, B. J. (2018). Meso-Cenozoic evolution of the southwestern Chukchi Borderland, Arctic Ocean. *Marine and Petroleum Geology*, 95, 100–109. <https://doi.org/10.1016/j.marpetgeo.2018.04.014>
- Ivanov, V. V., Shapiro, G. I., Huthnance, J. M., Aleynik, D. L., & Golovin, P. N. (2004). Cascades of dense water around the world ocean. *Progress in Oceanography*, 60(1), 47–98. <https://doi.org/10.1016/j.pcean.2003.12.002>
- Jakobsson, M. (1999). First high-resolution chirp sonar profiles from the central Arctic Ocean reveal erosion of Lomonosov Ridge sediments. *Marine Geology*, 158(1), 111–123. [https://doi.org/10.1016/S0025-3227\(98\)00186-8](https://doi.org/10.1016/S0025-3227(98)00186-8)
- Jakobsson, M., Andreassen, K., Bjarnadóttir, L. R., Dove, D., Dowdeswell, J. A., England, J. H., et al. (2014). Arctic Ocean glacial history. *Quaternary Science Reviews*, 92, 40–67. <https://doi.org/10.1016/j.quascirev.2013.07.033>
- Jakobsson, M., Gardner, J. V., Vogt, P. R., Mayer, L. A., Armstrong, A., Backman, J., et al. (2005). Multibeam bathymetric and sediment profiler evidence for ice grounding on the Chukchi Borderland, Arctic Ocean. *Quaternary Research*, 63(2), 150–160. <https://doi.org/10.1016/j.yqres.2004.12.004>
- Jakobsson, M., Løvlie, R., Arnold, E. M., Backman, J., Polyak, L., Knutsen, J.-O., & Musatov, E. (2001). Pleistocene stratigraphy and paleoenvironmental variation from Lomonosov Ridge sediments, central Arctic Ocean. *Global and Planetary Change*, 31(1), 1–22. [https://doi.org/10.1016/S0921-8181\(01\)00110-2](https://doi.org/10.1016/S0921-8181(01)00110-2)
- Jakobsson, M., Mayer, L. A., Bringensparr, C., Castro, C. F., Mohammad, R., Johnson, P., et al. (2020). The international bathymetric chart of the Arctic Ocean version 4.0. *Scientific Data*, (1), 176.
- Jakobsson, M., Nilsson, J., Anderson, L., Backman, J., Björk, G., Cronin, T. M., et al. (2016). Evidence for an ice shelf covering the central Arctic Ocean during the penultimate glaciation. *Nature Communications*, 7, 10365.
- Jakobsson, M., Nilsson, J., O'Regan, M., Backman, J., Löwemark, L., Dowdeswell, J. A., et al. (2010). An Arctic Ocean ice shelf during MIS 6 constrained by new geophysical and geological data. *Quaternary Science Reviews*, 29(25), 3505–3517. <https://doi.org/10.1016/j.quascirev.2010.03.015>

- Jakobsson, M., Polyak, L., Edwards, M., Kleman, J., & Coakley, B. (2008). Glacial geomorphology of the Central Arctic Ocean: The Chukchi Borderland and the Lomonosov Ridge. *Earth Surface Processes and Landforms*, 33(4), 526–545. <https://doi.org/10.1002/esp.1667>
- Joe, Y. J., Polyak, L., Schreck, M., Niessen, F., Yoon, S. H., Kong, G. S., & Nam, S.-I. (2020). Late Quaternary depositional and glacial history of the Arliss Plateau off the East Siberian margin in the western Arctic Ocean. *Quaternary Science Reviews*, 228, 106099. <https://doi.org/10.1016/j.quascirev.2019.106099>
- Jokat, W. (2009). *The expedition of the research vessel "Polarstern" to the Arctic in 2008 (ARK-XXIII/3)*. Bremerhaven, Germany: Alfred Wegener Institute.
- Kim, J.-H., Hachikubo, A., Kida, M., Minami, H., Lee, D.-H., Jin, Y. K., et al. (2020). Upwarding gas source and postgenetic processes in the shallow sediments from the ARAON Mounds, Chukchi Sea. *Journal of Natural Gas Science and Engineering*, 76, 103223.
- Kirkbride, M. P. (2002). 6—Processes of glacial transportation. In J. Menzies (Ed.), *Modern and past glacial environments* (pp. 147–169). Oxford: Butterworth-Heinemann. <https://doi.org/10.1016/B978-075064226-2/50009-X>
- Klages, J. P., Kuhn, G., Graham, A. G. C., Hillenbrand, C.-D., Smith, J. A., Nitsche, F. O., et al. (2015). Palaeo-ice stream pathways and retreat style in the easternmost Amundsen Sea Embayment, West Antarctica, revealed by combined multibeam bathymetric and seismic data. *Geomorphology*, 245, 207–222.
- Laberg, J. S., Vorren, T. O., Dowdeswell, J. A., Kenyon, N. H., & Taylor, J. (2000). The Andøya Slide and the Andøya Canyon, north-eastern Norwegian–Greenland Sea. *Marine Geology*, 162(2–4), 259–275. [https://doi.org/10.1016/S0025-3227\(99\)00087-0](https://doi.org/10.1016/S0025-3227(99)00087-0)
- LeDoux, C. M., Hulbe, C. L., Forbes, M. P., Scambos, T. A., & Alley, K. (2017). Structural provinces of the Ross Ice Shelf, Antarctica. *Annals of Glaciology*, 58(75 Pt 1), 88–98. <https://doi.org/10.1017/aog.2017.24>
- Lisiecki, L. E., & Raymo, M. E. (2005). A Pliocene–Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography*, 20, PA1003. <https://doi.org/10.1029/2004PA001071>
- Lowe, A., & Anderson, J. (2002). Reconstruction of the West Antarctic ice sheet in Pine Island Bay during the Last Glacial Maximum and its subsequent retreat history. *Quaternary Science Reviews*, 21(16–17), 1879–1897. [https://doi.org/10.1016/S0277-3791\(02\)00006-9](https://doi.org/10.1016/S0277-3791(02)00006-9)
- Lowe, A. L., & Anderson, J. B. (2003). Evidence for abundant subglacial meltwater beneath the paleo-ice sheet in Pine Island Bay, Antarctica. *Journal of Glaciology*, 49(164), 125–138.
- MacGregor, J. A., Fahnestock, M. A., Catania, G. A., Aschwanden, A., Clow, G. D., Colgan, W. T., et al. (2016). A synthesis of the basal thermal state of the Greenland Ice Sheet. *Journal of Geophysical Research: Earth Surface*, 121, 1328–1350. <https://doi.org/10.1002/2015JF003803>
- Matthiessen, J., Niessen, F., Stein, R., & Naafs, B. D. A. (2010). Pleistocene glacial marine sedimentary environments at the eastern Mendeleev Ridge, Arctic Ocean. *Polarforschung*, 79(2), 123–137.
- Menzies, J., & Shilts, B. W. (2002). 8—Subglacial environments. In J. Menzies (Ed.), *Modern and past glacial environments* (pp. 183–278). Oxford: Butterworth-Heinemann. <https://doi.org/10.1016/B978-075064226-2/50011-8>
- Menzies, J., & van der Meer, J. J. M. (2018). In J. Menzies & J. J. M. van der Meer (Eds.), *Past glacial environments*. (2nd ed.). Hoboken, NJ: Elsevier.
- Miramontes, E., Cattaneo, A., Jouet, G., Théreau, E., Thomas, Y., Rovere, M., et al. (2016). The Pianosa Contourite Depositional System (Northern Tyrrhenian Sea): Drift morphology and Plio-Quaternary stratigraphic evolution. *Marine Geology*, 378, 20–42. <https://doi.org/10.1016/j.margeo.2015.11.004>
- Montelli, A., Dowdeswell, J. A., Ottesen, D., & Johansen, S. E. (2018). Architecture and sedimentary processes on the mid-Norwegian continental slope: A 2.7 Myr record from extensive seismic evidence. *Quaternary Science Reviews*, 192, 185–207. <https://doi.org/10.1016/j.quascirev.2018.05.034>
- Moore, D. G. (1964). Shear strength and related properties of sediments from experimental mohole (Guadalupe site). *Journal of Geophysical Research*, 69(20), 4271–4291.
- Newton, A. M. W., Knutz, P. C., Huuse, M., Gannon, P., Brocklehurst, S. H., Clausen, O. R., & Gong, Y. (2017). Ice stream reorganization and glacial retreat on the northwest Greenland shelf. *Geophysical Research Letters*, 44, 7826–7835. <https://doi.org/10.1002/2017GL073690>
- Niessen, F., Hong, J. K., Hegewald, A., Matthiessen, J., Stein, R., Kim, H., et al. (2013). Repeated Pleistocene glaciation of the East Siberian continental margin. *Nature Geoscience*, 6(10), 842–846. <https://doi.org/10.1038/ngeo1904>
- Noormets, R., Dowdeswell, J. A., Larter, R. D., Ó Cofaigh, C., & Evans, J. (2009). Morphology of the upper continental slope in the Bellingshausen and Amundsen Seas—Implications for sedimentary processes at the shelf edge of West Antarctica. *Marine Geology*, 258(1), 100–114. <https://doi.org/10.1016/j.margeo.2008.11.011>
- Ó Cofaigh, C., Dowdeswell, J. A., Allen, C. S., Hiemstra, J. F., Pudsey, C. J., Evans, J., & Evans, D. J. A. (2005). Flow dynamics and till genesis associated with a marine-based Antarctic palaeo-ice stream. *Quaternary Science Reviews*, 24(5), 709–740. <https://doi.org/10.1016/j.quascirev.2004.10.006>
- Ó Cofaigh, C., Dowdeswell, J. A., Evans, J., Kenyon, N. H., Taylor, J., Mienert, J., & Wilken, M. (2004). Timing and significance of glacially influenced mass-wasting in the submarine channels of the Greenland Basin. *Marine Geology*, 207(1), 39–54. <https://doi.org/10.1016/j.margeo.2004.02.009>
- Ó Cofaigh, C., Hogan, K. A., Jennings, A. E., Callard, S. L., Dowdeswell, J. A., Noormets, R., & Evans, J. (2018). The role of meltwater in high-latitude trough-mouth fan development: The Disko Trough-Mouth Fan, West Greenland. *Marine Geology*, 402, 17–32. <https://doi.org/10.1016/j.margeo.2018.02.001>
- Ó Cofaigh, C., Taylor, J., Dowdeswell, J. A., & Pudsey, C. J. (2003). Palaeo-ice streams, trough mouth fans and high-latitude continental slope sedimentation. *Boreas*, 32(1), 37–55.
- O'Grady, D. B., & Syvitski, J. P. (2002). Large-scale morphology of Arctic continental slopes: The influence of sediment delivery on slope form. In J. A. Dowdeswell & C. Ó Cofaigh (Eds.), *Special Publication—Geological Society of London* (Vol. 203, pp. 11–32). London: Geological Society of London.
- O'Regan, M., Backman, J., Barrientos, N., Cronin, T. M., Gemery, L., Kirchner, N., et al. (2017). The De Long Trough: A newly discovered glacial trough on the East Siberian continental margin. *Climate of the Past*, 13(9), 1269–1284. <https://doi.org/10.5194/cp-13-1269-2017>
- Ottesen, D., & Dowdeswell, J. A. (2006). Assemblages of submarine landforms produced by tidewater glaciers in Svalbard. *Journal of Geophysical Research*, 111, F01016. <https://doi.org/10.1029/2005JF000330>
- Ottesen, D., & Dowdeswell, J. A. (2009). An inter-ice-stream glaciated margin: Submarine landforms and a geomorphic model based on marine–geophysical data from Svalbard. *The Geological Society of America Bulletin*, 121(11–12), 1647–1665.
- Ottesen, D., Rise, L., Knies, J., Olsen, L., & Henriksen, S. (2005). The Vestfjorden-Trænadjupet palaeo-ice stream drainage system, mid-Norwegian continental shelf. *Marine Geology*, 218(1), 175–189. <https://doi.org/10.1016/j.margeo.2005.03.001>
- Park, K., Kim, J.-H., Asahi, H., Polyak, L., Khim, B.-K., Schreck, M., et al. (2020). Cyclostratigraphic age constraining for Quaternary sediments in the Makarov Basin of the western Arctic Ocean using manganese variability. *Quaternary Geochronology*, 55, 101021.

- Paull, C. K., Normark, W. R., Ussler, W., Caress, D. W., & Keaten, R. (2008). Association among active seafloor deformation, mound formation, and gas hydrate growth and accumulation within the seafloor of the Santa Monica Basin, offshore California. *Marine Geology*, 250(3), 258–275.
- Piper, D. J. W., Deptuck, M., Mosher, D., Hughes Clarke, J., Migeon, S., Prather, B., et al. (2012). Erosional and depositional features of glacial meltwater discharges on the eastern Canadian continental margin. In B. E. Prather, M. E. Deptuck, D. Mohrig, B. van Hoon, & R. Wynn (Eds.), *Applications of the principles of seismic geomorphology to continental slope and base-of-slope systems: Case studies from seafloor and near-seafloor analogues*. Society for Sedimentary Geology (SEPM), special publications (Vol. 99, p. 80). Tulsa, OK: SEPM.
- Piper, D. J. W., & Normark, W. R. (2009). Processes that initiate turbidity currents and their influence on turbidites: A marine geology perspective. *Journal of Sedimentary Research*, 79(6), 347–362. <https://doi.org/10.2110/jsr.2009.046>
- Polyak, L., Best, K. M., Crawford, K. A., Council, E. A., & St-Onge, G. (2013). Quaternary history of sea ice in the western Arctic Ocean based on foraminifera. *Quaternary Science Reviews*, 79, 145–156. <https://doi.org/10.1016/j.quascirev.2012.12.018>
- Polyak, L., Darby, D. A., Bischof, J. F., & Jakobsson, M. (2007). Stratigraphic constraints on late Pleistocene glacial erosion and deglaciation of the Chukchi margin, Arctic Ocean. *Quaternary Research*, 67(2), 234–245. <https://doi.org/10.1016/j.yqres.2006.08.001>
- Polyak, L., Edwards, M. H., Coakley, B. J., & Jakobsson, M. (2001). Ice shelves in the Pleistocene Arctic Ocean inferred from glaciogenic deep-sea bedforms. *Nature*, 410, 453–457.
- Polyak, L., Forman, S. L., Herlihy, F. A., Ivanov, G., & Krinitsky, P. (1997). Late Weichselian deglacial history of the Svyataya (Saint) Anna Trough, northern Kara Sea, Arctic Russia. *Marine Geology*, 143(1), 169–188. [https://doi.org/10.1016/S0025-3227\(97\)00096-0](https://doi.org/10.1016/S0025-3227(97)00096-0)
- Pope, E. L., Talling, P. J., & Ó Cofaigh, C. (2018). The relationship between ice sheets and submarine mass movements in the Nordic Seas during the Quaternary. *Earth-Science Reviews*, 178, 208–256. <https://doi.org/10.1016/j.earscirev.2018.01.007>
- Rebesco, M., & Camerlenghi, A. (2008). Late Pliocene margin development and mega debris flow deposits on the Antarctic continental margins: Evidence of the onset of the modern Antarctic Ice Sheet? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 260(1), 149–167. <https://doi.org/10.1016/j.palaeo.2007.08.009>
- Rebesco, M., Özmaral, A., Urgeles, R., Accettella, D., Lucchi, R. G., Rütther, D., et al. (2016). Evolution of a high-latitude sediment drift inside a glacially-carved trough based on high-resolution seismic stratigraphy (Kveithola, NW Barents Sea). *Quaternary Science Reviews*, 147, 178–193. <https://doi.org/10.1016/j.quascirev.2016.02.007>
- Rignot, E., Mouginot, J., & Scheuchl, B. (2011). Ice flow of the Antarctic ice sheet. *Science*, 333(6048), 1427–1430.
- Rydningen, T. A., Laberg, J. S., & Kolstad, V. (2015). Seabed morphology and sedimentary processes on high-gradient trough mouth fans offshore Troms, northern Norway. *Geomorphology*, 246, 205–219. <https://doi.org/10.1016/j.geomorph.2015.06.007>
- Schomacker, A., Kjær, K. H., & Krüger, J. (2010). 8: Subglacial environments, sediments and landforms at the margins of Mýrdalsjökull. In A. Schomacker, J. Krüger, & K. H. Kjær (Eds.), *The Mýrdalsjökull Ice Cap, Iceland. Glacial processes, sediments and landforms on an active volcano* (Vol. 13, pp. 127–144). Hoboken, NJ: Elsevier. [https://doi.org/10.1016/S1571-0866\(09\)01308-6](https://doi.org/10.1016/S1571-0866(09)01308-6)
- Schreck, M., Nam, S.-I., Polyak, L., Vogt, C., Kong, G.-S., Stein, R., et al. (2018). Improved Pleistocene sediment stratigraphy and paleoenvironmental implications for the western Arctic Ocean off the East Siberian and Chukchi margins. *Arktos*, 4(1), 21. <https://doi.org/10.1007/s41063-018-0057-8>
- Shaver, R., & Hunkins, K. (1964). Arctic Ocean geophysical studies: Chukchi Cap and Chukchi abyssal plain. *Deep Sea Research and Oceanographic Abstracts*, 11(6), 905–916. [https://doi.org/10.1016/0011-7471\(64\)90340-7](https://doi.org/10.1016/0011-7471(64)90340-7)
- Shipp, S., Anderson, J., & Domack, E. (1999). Late Pleistocene–Holocene retreat of the West Antarctic Ice-Sheet system in the Ross Sea: Part 1—Geophysical results. *The Geological Society of America Bulletin*, 111(10), 1486–1516.
- Solheim, A., Faleide, J. I., Andersen, E. S., Elverhøi, A., Forsberg, C. F., Vanneste, K., et al. (1998). Late Cenozoic seismic stratigraphy and glacial geological development of the East Greenland and Svalbard–Barents Sea continental margins. *Quaternary Science Reviews*, 17(1), 155–184. [https://doi.org/10.1016/S0277-3791\(97\)00068-1](https://doi.org/10.1016/S0277-3791(97)00068-1)
- Spagnolo, M., Clark, C. D., Ely, J. C., Stokes, C. R., Anderson, J. B., Andreassen, K., et al. (2014). Size, shape and spatial arrangement of mega-scale glacial lineations from a large and diverse dataset. *Earth Surface Processes and Landforms*, 39(11), 1432–1448. <https://doi.org/10.1002/esp.3532>
- Stärz, M., Gong, X., Stein, R., Darby, D. A., Kauker, F., & Lohmann, G. (2012). Glacial shortcut of Arctic sea-ice transport. *Earth and Planetary Science Letters*, 357–358, 257–267. <https://doi.org/10.1016/j.epsl.2012.09.033>
- Stein, R., Matthießen, J., Niessen, F., Krylov, A., Nam, S., & Bazhenova, E. (2010). Toward a better (litho-) stratigraphy and reconstruction of Quaternary paleoenvironment in the Amerasian Basin (Arctic Ocean). *Polarforschung*, 79(2), 97–121.
- Stewart, M. A., & Lonergan, L. (2011). Seven glacial cycles in the middle-late Pleistocene of northwest Europe: Geomorphic evidence from buried tunnel valleys. *Geology*, 39(3), 283–286. <https://doi.org/10.1130/G31631.1>
- SWERUS Scientific Party. (2016). *Cruise report SWERUS-C3 Leg 2*.
- Taylor, J., Dowdeswell, J. A., Kenyon, N. H., & Ó Cofaigh, C. (2002). Late Quaternary architecture of trough-mouth fans: Debris flows and suspended sediments on the Norwegian margin. In J. A. Dowdeswell & C. Ó Cofaigh (Eds.), *Glacier-influenced sedimentation on high-latitude continental margins* (pp. 55–71). London: Geological Society of London. <https://doi.org/10.1144/GSL.SP.2002.203.01.04>
- Todd, B. J., Valentine, P. C., Longva, O., & Shaw, J. (2007). Glacial landforms on German Bank, Scotian Shelf: Evidence for Late Wisconsinan ice-sheet dynamics and implications for the formation of De Geer moraines. *Boreas*, 36(2), 148–169. <https://doi.org/10.1111/j.1502-3885.2007.tb01189.x>
- Veeken, P. C. H., & van Moerkerken, B. (2013). *Seismic stratigraphy and depositional facies models*. Houten, The Netherlands: EAGE Publications.
- Vorren, T., & Laberg, J. S. (1997). Trough mouth fans—Palaeoclimate and ice-sheet monitors. *Quaternary Science Reviews*, 16(8), 865–881.
- Waage, M., Portnov, A., Serov, P., Bünz, S., Waghorn, K. A., Vadakkupuliyambatta, S., et al. (2019). Geological controls on fluid flow and gas hydrate pingo development on the Barents Sea margin. *Geochemistry, Geophysics, Geosystems*, 20, 630–650. <https://doi.org/10.1029/2018GC007930>
- Wang, R., Xiao, W., März, C., & Li, Q. (2013). Late Quaternary paleoenvironmental changes revealed by multi-proxy records from the Chukchi Abyssal Plain, western Arctic Ocean. *Global and Planetary Change*, 108, 100–118. <https://doi.org/10.1016/j.gloplacha.2013.05.017>
- Wellner, J. S., Heroy, D. C., & Anderson, J. B. (2006). The death mask of the Antarctic ice sheet: Comparison of glacial geomorphic features across the continental shelf. *Geomorphology*, 75(1), 157–171. <https://doi.org/10.1016/j.geomorph.2005.05.015>
- Wellner, J. S., Lowe, A. L., Shipp, S. S., & Anderson, J. B. (2001). Distribution of glacial geomorphic features on the Antarctic continental shelf and correlation with substrate: Implications for ice behavior. *Journal of Glaciology*, 47(158), 397–411. <https://doi.org/10.3189/172756501781832043>
- Wille, P. (2005). *Sound images of the ocean: In research and monitoring* (Vol. 1). New York: Springer Science & Business Media.

- Winsborrow, M. C. M., Clark, C. D., & Stokes, C. R. (2010). What controls the location of ice streams? *Earth-Science Reviews*, 103(1–2), 45–59. <https://doi.org/10.1016/j.earscirev.2010.07.003>
- Xiao, W., Polyak, L., Wang, R., Löwemark, L., Mei, J., You, D., et al. (2020). Middle to Late Pleistocene Arctic paleoceanographic changes based on sedimentary records from Mendeleev Ridge and Makarov Basin. *Quaternary Science Reviews*, 228, 106105. <https://doi.org/10.1016/j.quascirev.2019.106105>
- Zaragosi, S., Auffret, G. A., Faugères, J.-C., Garlan, T., Pujol, C., & Cortijo, E. (2000). Physiography and recent sediment distribution of the Celtic Deep-Sea Fan, Bay of Biscay. *Marine Geology*, 169(1), 207–237.