



Full length article

Immune gene discovery in the crucian carp *Carassius auratus*

Jae-Sung Rhee^a, Chang-Bum Jeong^b, Duck-Hyun Kim^c, Il-Chan Kim^d, Yong Sung Lee^e,
Chulwoo Lee^{f,**}, Jae-Seong Lee^{g,*}

^a Research Institute for Natural Sciences, Hanyang University, Seoul 133-791, South Korea

^b Department of Chemistry, College of Natural Sciences, Hanyang University, Seoul 133-791, South Korea

^c Department of Life Science, College of Convergence, Sangmyung University, Seoul 110-743, South Korea

^d Division of Life Sciences, Korea Polar Research Institute, Incheon 406-840, South Korea

^e Graduate School of Biomedical Science and Engineering, Hanyang University, Seoul 133-791, South Korea

^f Research Strategy & Planning Division, National Institute of Environmental Research, Incheon 404-708, South Korea

^g Department of Biological Sciences, College of Natural Sciences, Sungkyunkwan University, Suwon 440-746, South Korea

ARTICLE INFO

Article history:

Received 7 August 2013

Received in revised form

6 November 2013

Accepted 7 November 2013

Available online 26 November 2013

Keywords:

Crucian carp

Carassius auratus

Pyrosequencing

Immune-relevant genes

ABSTRACT

The crucian carp *Carassius auratus* (Cyprinidae) is one of the important fish species in aquaculture. Although the crucian carp has several economic benefits, their immune system and gene information have not been investigated in depth as yet. Here, we performed the transcriptome analysis of *C. auratus* using the pyrosequencing method and selected several immune-related genes. Of unigenes obtained in this species, we identified a number of immune system-related genes (e.g. adhesive protein, antimicrobial protein, apoptosis- and cell cycle-related protein, cellular defense effector, immune regulator, pattern recognition protein, protease, protease inhibitor, reduction/oxidation-related protein, signal transduction-related protein and stress protein) that are potentially useful for studies on fish immunity. To be of public and practical use, we designed primer pairs of each gene from the crucian carp for real-time RT-PCR application and tested the amplicon identity of entire gene sets with the total RNA sample. For comparative analysis, we measured tissue-preferential transcript profiles of selected genes. This study will be helpful to extend our knowledge on the immune system of the crucian carp in comparative aspects and to develop the crucian carp as a potential model organism for aquatic quality monitoring in fish farming.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

The crucian carp *Carassius auratus* is a member of the family Cyprinidae (Teleostei). These fishes are world-wide distributed in freshwater systems such as river, lake, and pond. Previously, *C. auratus* was highlighted as one of most important commercial species in aquaculture including several marine fish and crustaceans as their large production, for their excellent growth traits, good taste, and suitability in culture systems (FAO; Food and Agriculture Organization of the United Nations). Although the crucian carp is highly susceptible to bacterial and pathogen challenges, exploration of immunity at the molecular level is dominated by several aquaculture model animals by diverse experimental approaches with little attention to the crucian carp *C. auratus*. To

date, in crucian carp, several immunity-relevant factors such as interferon regulatory factor 7 [63], *CD8α* [46], interferon-stimulated gene ISG15 [64], *GATA3* [49], MHC class I and β -2 microglobulin [52], and *IRF9* [45] were cloned and characterized but have not been reported as yet for the identification of extensive immune genes and their employment for immunity. Therefore, investigation of key immune components and characterization of gene/protein expression of immunity-relevant genes would have priority in the immune research of crucian carp.

Molecular genomic and proteomic approaches are able to provide a better understanding on the intracellular mechanism on the immune system [14,47]. Recently, several Next Generation Sequencing (NGS) technologies enable us to obtain the massive complementary DNA (cDNA) or genomic DNA (gDNA) information, and thus to date the extensive sequence information of important immune-relevant genes has been identified in several fish species and employed for further understanding of immune system [2,14,47,55]. On the usefulness of the crucian carp as a potential model species for aquaculture monitoring, we sequenced transcriptomes for the gene/protein information-based application to

* Corresponding author. Department of Biological Sciences, College of Natural Sciences, Sungkyunkwan University, Suwon 440-746, South Korea. Tel.: +82 2 447 2382.

** Corresponding author.

E-mail addresses: leecwoo@korea.kr (C. Lee), r_marmoratus@hanmail.net (J.-S. Lee).

detect early molecular biomarker genes of the disease induction or pollution in crucian carp.

In this study, we obtained expressed cDNA information by pyrosequencing in the crucian carp, *C. auratus*. Our results are the first report on the extensive identification of various immune mechanism related genes in this species. To employ transcriptional profiling for aquaculture monitoring, we designed primer sets for entire immune-relevant genes of the crucian carp and tested amplicon identity for public and practical use in immunity research. Finally, we supposed that these results would be useful to obtain a better understanding of the mode of action of disease induction by pathogens in the crucian carp, *C. auratus*.

2. Materials and methods

2.1. Fish

The crucian carp *C. auratus* were reared and maintained at the National Institute of Environmental Research (Incheon, South Korea). The fish were maintained at 25 °C with 12 h/12 h light/darkness. The automated water changing system was set for constant flow-through and water quality (pH, salinity, and temperature) was recorded using various instruments (5.71 ± 0.19 mg O₂/L). Experimental fish were anaesthetized on ice and sacrificed by decapitation.

2.2. Total RNA extraction

The main objective of this study was the mining of expressed transcripts that provide a potentially immune-relevant gene pool of the crucian carp for public use in diverse immune research. Therefore, to enrich transcriptome information, we sampled different total RNA pools from immunized or non-immunized tissues. In particular tissues collected from brain, eye, gill, heart, head kidney, intestine, kidney, liver, muscle, skin, spleen, and stomach of 10 fish at the adult stage (≈ 8 month) of both sexes. Each pooled tissues were homogenized in 3 volumes of TRIZOL[®] reagent (Molecular Research Center, Inc., Cincinnati, OH) with a tissue grinder and stored at -80 °C until use. Total RNAs were isolated from each pooled tissues according to the manufacturer's instructions. DNA digestion was performed using DNase I (Sigma, St. Louis, Mo). After DNase treatment, the total RNAs were purified using RNeasy Mini Spin Column (Qiagen), and were quantified by absorption of light at 230, 260, and 280 nm (A230/260, A260/280) using a spectrophotometer (Ultrospec 2100 pro, Amersham Bioscience). To check the genomic DNA contamination, we loaded the total RNAs in a 1% agarose gel which contained ethidium bromide (EtBr) and visualized on a UV transilluminator (Wealtec Corp., Sparks, NV). Also, to verify the total RNA quality, we loaded the total RNAs in a 1% formaldehyde/agarose gel with EtBr staining and checked the 18/28S ribosomal RNAs integrity. After RNA quality was determined, single-strand cDNA was synthesized from total RNA using an oligo(dT)₂₀ primer for reverse transcription (SuperScript[™] III RT kit, Invitrogen, Carlsbad, CA).

2.3. cDNA library preparation

cDNA library was constructed according to the manufacturer's instructions (Roche Applied Science, Genome Sequencer 20[™] System). Briefly, cDNA was fractionated into smaller fragments (300–500 base pairs) that were subsequently blunt-ended using Fragment End Polishing kit supplied by manufacture (Roche Applied Science). Short Adaptors (Two adaptors, A and B, were provided in the GS-20 Library Preparation Kit) were then ligated onto the ends of the fragments. These adaptors were provided priming for both

amplification and sequencing of the sample-library fragments. Adaptor B contained a 5'-biotin tag that enables immobilization of the library onto streptavidin-coated beads. After fill-in reaction for nicks, the non-biotinylated strand was used to isolate a single-stranded template DNA (sstDNA) library. The sstDNA library was assessed for its quality and the optimal amount (DNA copies per bead) needed for emulsion PCR (emPCR) was determined by titration.

2.4. Pyrosequencing

Expressed cDNA sequencing was performed with GS 20 sequencer. Constructed sstDNA library was clonally amplified via emPCR, thereby generating millions of copies of templates per bead. The DNA beads were then distributed into picolitre-sized wells on a fiber-optic slide (PicoTiter-Plate[™]) along with a mixture of smaller beads coated with the enzymes required for the pyrosequencing reaction including the firefly enzyme luciferase. The four DNA nucleotides were then flushed sequentially over the plate. Light signals released upon base incorporation were captured by a CCD camera, and the sequence of bases incorporated per well was stored as a read.

2.5. Assembly, gene annotation, and GO analysis

For assembly, we used the Newbler Assembler software (454 Life Sciences, Roche Diagnostics company) in the assembly stage 1, and the assembled EST translations were aligned with the GenBank NR (non-redundant; including all GenBank, EMBL, DDBJ, and PDB sequence except EST, STS, GSS, or HTGS) amino acid sequence database using BLASTx. The aligned data were arranged according to read length, gene annotation, GenBank number, *E*-value, species, and its accession number. All the bacterial clones were eliminated from the annotated genes. The Gene ontology (GO) functional annotations were assigned by Blast2GO automated sequence annotation tool ([11]; <http://www.blast2go.org>). Three main categories for biological process, cellular component, and molecular function were obtained by using default parameters. Assembly, gene annotation, and GO analysis were performed at the National Instrumentation Center for Environmental Management (NICEM), Seoul National University (Seoul, South Korea).

2.6. Immune-relevant gene mining

In each section of GO terms, immune-relevant gene mining was performed manually within the expected value threshold of 1.00E-05 that has potential immune-relevant function. Selected genes were arranged with read length, gene annotation, GenBank number, *E*-value, and species with its accession number.

2.7. Primer design and real-time RT-PCR

Primers were designed after comparing exon/intron boundary to genomic DNA using GENRUNNER software (Hastings Software, Inc. N. Y. USA) and confirmed by Primer 3 program (Whitehead Institute/MIT center for Genome Research). To determine the amplicon identity, all the PCR products were cloned into pCR2.1 TA vector, and sequenced with an ABI 3700 DNA analyzer (Bionics Co., Seoul, South Korea). To analyze transcriptional expression patterns in different tissues of the crucian carp, several immune-relevant genes were collected from the assembled EST database, and their transcript abundances were analyzed with real-time RT-PCR. Template cDNA was made by pooled total RNA of each tissue from 10 adult fish of both sexes. Each reaction included 1 μ l of cDNA and 0.2 μ M primer (real-time RT-F/R or 18S rRNA RT-F/R). Optimized

conditions were transferred according to the following CFX96™ real-time PCR protocol (Bio-Rad, Hercules, CA, USA). Reaction conditions were as follows; 95 °C/3 min; 40 cycles of 95 °C/30 s, 55 °C/30 s, 72 °C/30 s. To confirm the proper amplification of specific products, cycles were continued to check the melting curve under the following conditions; 95 °C/1 min, 55 °C/1 min, and 80 cycles of 55 °C/10 s with 0.5 °C increase per cycle. SYBR® Green (Molecular Probes Inc., Invitrogen) was used to detect specific amplified products. Amplification and detection of SYBR® Green-labeled products were performed using CFX96™ real-time PCR system (Bio-Rad, Hercules, CA, USA). Data from each experiment were expressed relative to expression levels of the 18S rRNA gene to normalize the expression levels between samples. All the experiments were done in triplicate. Data were collected as threshold cycle (C_T) values (PCR cycle number where fluorescence was detected above a threshold and decreased linearly with increasing input target quantity), and used to calculate ΔC_T values of each sample. The fold change in the relative gene expression was calculated by the $2^{-\Delta\Delta C_T}$ method [27].

2.8. Statistical analysis

Data were expressed as mean \pm standard deviation (S.D.). Significant differences were analyzed using one-way and/or multiple-comparison ANOVA followed by Tukey's post hoc test. $P < 0.05$ was considered significant. SPSS ver. 17.0 (SPSS Inc., Chicago IL, USA) software package was used for statistical analysis.

3. Results and discussion

3.1. Overview of the crucian carp ESTs

Overall pyrosequencing result on transcriptome of the crucian carp was summarized in Table 1. As raw read data, we obtained 221,542 reads, and all the reads were trimmed and cleaned with several bioinformatics tools. Repetitive and low quality reads were

filtered out from the entire raw read set for non-redundant consensus. After the cleaning process, 19,492 contigs and 58,743 singletons were aligned, respectively. After all the transcripts were compared with the GenBank NR amino acid sequence database using BLASTx, those assembled transcripts with E -values $\leq 1.00E-05$ were used to obtain reliable annotation results in this study. The average read length of those was 106 bp in contig and 84 bp in singleton reads, respectively. The largest contig length was 1012 bp and the largest singleton length was 191 bp, respectively. Regarding sequencing quality and average read lengths of views in using the GS-20 platform, overall pyrosequencing results of the crucian carp were similar to previous results using the bacterium *Mycoplasma genitalium* (110 bp) [31], the hermaphroditic fish *Kryptolebias marmoratus* transcripts (116–129 bp) [36], and the rockshell *Thais clavigera* (114 bp) [37] that were sequenced by the GS-20 pyrosequencing method. To increase valid nomenclatures, entire reads were sorted out with the cut-off value more than 200 bp with E -value $\leq 1.00E-05$. Subsequently, selected reads were analyzing their Gene Ontology (GO) to find specific gene function of their own as well as their immune-relevant role.

3.2. Annotation and functional classification for immune-relevant gene mining

Of the 78,235 contig and singleton cDNA fragments, 10,532 reads (13%) were aligned with valid similarities in the NCBI database (Table 1). In 10,532 reads, 7625 genes (72%) were matched with those of fish species. Those genes were classified according to their functions into each ontology class such as molecular function, cellular process, or biological process in Blast2GO analysis. Briefly, the vast majority of genes were involved in “binding” (49%), and “catalytic activity” subcategories (28%) in the molecular function category (Fig. 1A). In the biological process class, most genes were categorized in “cellular processes” (15%) and “metabolic processes” (15%) (Fig. 1B). The “immune system process” subcategory was occupied by 3% of the biological process category. In the cellular component class, most of genes were related to “cell” (22%) and “cell part” (22%) and followed by the “organelle” (16%) subcategory (Fig. 1C). Based on functional analysis, we identified immune-relevant genes from each subcategory as many of them were incorporated into different categories for coordination of diverse functions with other genes. Finally, 77 immune system-related sequences corresponding to adhesive protein, antimicrobial protein, apoptosis- and cell cycle-related protein, cellular defense effector, immune regulator, protease and protease inhibitor, reduction/oxidation-related protein, signal transduction-related protein, stress protein, and acute phase response/inflammation-related genes were identified in the EST database for further work (Table 2). Taken together, pyrosequencing-based GO profiling is of value in the identification of immune-relevant genes, indicating that this kind of approach would be available to better understand the functional role of certain gene sets or signaling pathways.

3.3. Immune-relevant genes in the crucian carp

To provide public and practical use of immune-relevant genes, we designed primer pairs for the entire immune gene set identified in the crucian carp (Table 3) and tested their amplicon identity with electrophoresis (Fig. 2). Subsequently, we measured tissue-preferential transcript profiles of selected genes that were confirmed in other teleosts to validate the usefulness of gene-specific applications for further immune-relevant work in the crucian carp.

Table 1
Overview of pyrosequencing and assembly result.

# EST sequences	221,542
Sequence cleaning process ^a	
# valid reads	203,281
# trimmed reads of valid process	5947
# trashed reads ^b	12,314
Lucy cleaning process	
# valid reads	193,553
# trashed reads	9728
Assembly Stage ^c	
# contigs (2+) ^d	19,492
average read length (bp)	106
largest read length (bp)	1012
# singletons ^e	58,743
average read length (bp)	84
largest read length (bp)	191
# transcripts ^f	78,235
# hits ^g	10,532
% with hits ^h	13

^a Minimum length: 60 base pair.

^b Trimmed reads composed of short reads (7341), low quality reads (1657), short quality reads (3241), and dust reads (75).

^c Assembly was performed using the Assembly Software (Newbler Assembler Software) of the Genome Sequencer 20 Software Version 1.0.53.

^d Number of contigs with 2 or more sequences.

^e Number of contigs with 1 sequence.

^f Total number of transcripts including singletons.

^g Number of transcripts that have a BLASTX hit.

^h Percent of assembled transcripts that have a BLASTX hit.




Fig. 1. Categories of gene ontology for the crucian carp ESTs. (A) molecular function, (B) biological process, (C) cellular components.

3.3.1. C-type lectin

Lectin is a large glycoprotein family that is capable of binding sugar moieties of foreign pathogens with a carbohydrate recognition domain (CRD) [12]. Lectins are classified according to their secondary and/or tertiary structure as calnexin, C-, L-, P-, I-, R-, and S-type lectins. Of lectins, the C-type lectins as a large group of lectins are pattern-recognition proteins which are functionally important for pathogen recognition and immune regulation in animals [58]. In the crucian carp, the highest mRNA expression level of C-type lectin (*Ca-CTL*) was observed in liver, while moderate expression level in intestine, spleen, and muscle tissues (Fig. 3). Previously, Kondo et al. [21] reported that C-type lectin transcript was only observed in liver tissues of the Japanese flounder, *Paralichthys olivaceus*. In the orange-spotted grouper *Epinephelus coioides*, C-type lectin transcript was abundant in liver and skin [53]. Also, transcriptional abundance of C-type lectin was highest in liver but was lowest in kidney [62]. On the contrary, dissimilar transcriptional profiles were observed in several teleosts [42,60]. Specific immune-related function of C-type lectin is still

unclear in different tissues of teleosts as yet, although some hepatitis C viral glycoproteins are known to invade liver cells for infection via attachment to C-type lectins in vertebrates [17,35]. Moreover, an appropriate gene family backbone is not available, as each lectin family is composed of large complex subfamilies. In the case of C-type lectin, amino acid similarity is not high within teleosts (identity: <81% in BLASTx). For example, *Ca-CTL* amino acid showed a moderate similarity to Japanese flounder (*P. olivaceus*, 41%) and orange-spotted grouper (*E. coioides*, 46%). Based on various isotypes of C-type lectin and their different transcriptional expressions in tissues, each C-type lectin would have potentially specialized immune-related roles in different tissues, and subsequent comparative phylogenetic and functional studies will be highlighted in elucidating the role of teleosts' C-type lectin family.

3.3.2. CC chemokine

Chemokines are a large family of cytokines that are classified according to structural characteristics and conserved cysteine residues [65]. Chemokines play a pivotal role in both, the non-specific

Table 2
Immune-relevant genes annotated in the EST library from *C. auratus*.

Putative function	GenBank	Species (Accession no.)	E-value
Adhesive protein			
Coagulation factor II	JX477179	<i>Danio rerio</i> (NP_998555)	0
Coagulation factor VII	JX477180	<i>Ictalurus furcatus</i> (ADO28333)	4E-30
Coagulation factor X	JX477181	<i>Danio rerio</i> (NP_958870)	5E-125
Coagulation factor V	JX477182	<i>Danio rerio</i> (AAN71005)	2E-176
C-type lectin	JX477183	<i>Ctenopharyngodon idella</i> (ACS16044)	1E-140
Fibrinogen, α chain	JX477186	<i>Danio rerio</i> (NP_001181918)	3E-67
Fibrinogen, β polypeptide	JX477184	<i>Danio rerio</i> (NP_997939)	2E-168
Fibrinogen, γ polypeptide	JX477185	<i>Hypophthalmichthys molitrix</i> (ADF97606)	4E-77
F11 receptor	JX477187	<i>Danio rerio</i> (NP_001076451)	9E-30
Intelectin 3	JX477188	<i>Ctenopharyngodon idella</i> (AAY43357)	1E-53
CD59 glycoprotein	JX477189	<i>Danio rerio</i> (XP_003198001)	1E-35
Integral membrane protein 2Bb	JX477190	<i>Danio rerio</i> (NP_998141)	5E-64
Antimicrobial			
Liver-expressed antimicrobial peptide 2B	JX477191	<i>Oncorhynchus mykiss</i> (NP_001117937)	6E-18
Apoptosis and cell cycle			
Programmed cell death 10	JX477192	<i>Ictalurus punctatus</i> (NP_001187364)	4E-37
Calmodulin	JX477193	<i>Homo sapiens</i> (AAH07965)	1E-66
Complement c1q, tumor necrosis factor-related protein	JX477194	<i>Danio rerio</i> (XP_001919564)	2E-41
Myeloid cell leukemia sequence 1B	JX477195	<i>Danio rerio</i> (AAL32470)	5E-41
Cellular defense effector			
α -1-microglobulin	JX477196	<i>Danio rerio</i> (NP_957412)	4E-119
α -2-macroglobulin	JX477198	<i>Cyprinus carpio</i> (ABC47740)	1E-99
β -2-microglobulin	JX477197	<i>Labeobarbus intermedius</i> (CAD44964)	6E-44
MHC class I antigen	JX477199	<i>Cyprinus carpio</i> (AF038426)	4E-23
Eukaryotic translation initiation factor 3 subunit M, dendritic cell protein	JX477200	<i>Danio rerio</i> (NP_001019906)	7E-47
Leukocyte cell-derived chemotaxin 2	JX477201	<i>Danio rerio</i> (NP_001041520)	3E-78
Chemokine CCL-c25s	JX477202	<i>Danio rerio</i> (BAF98242)	4E-23
Immune regulator			
Kininogen 1	JX477203	<i>Danio rerio</i> (NP_001005981)	1E-91
Secreted immunoglobulin domain 4	JX477204	<i>Danio rerio</i> (NP_001029354)	4E-107
Carboxypeptidase B2	JX477206	<i>Danio rerio</i> (NP_001018539)	4E-48
Signal sequence receptor, γ	JX477207	<i>Danio rerio</i> (NP_956347)	2E-67
Translocon-associated protein subunit γ			
Fetuin-A, α -2-HS-glycoprotein	JX477208	<i>Danio rerio</i> (NP_997787)	3E-35
Basic transcription factor 3	JX477209	<i>Danio rerio</i> (NP_001070026)	1E-39
Proteases and protease inhibitor			
Inter- α (globulin) inhibitor H3	JX477210	<i>Danio rerio</i> (NP_001018424)	1E-96
Inter- α -trypsin inhibitor heavy chain H4 precursor	JX477211	<i>Danio rerio</i> (NP_001020335)	3E-48
Serine (or cysteine) proteinase inhibitor, clade C (antithrombin)	JX477212	<i>Danio rerio</i> (NP_878283)	2E-103
Serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor)	JX477214	<i>Danio rerio</i> (CAQ14779)	7E-40
Serine (or cysteine) proteinase inhibitor, clade A (α -1 antiproteinase), member 1	JX477215	<i>Danio rerio</i> (AAH95615)	1E-60
Serpin peptidase inhibitor, clade A (α -1 antiproteinase), member 7	JX477213	<i>Danio rerio</i> (AAI53542)	4E-100
Serpin peptidase inhibitor, clade F, member 2B	JX477217	<i>Danio rerio</i> (NP_001073479)	3E-57
C1 inhibitor precursor	JX477216	<i>Danio rerio</i> (NP_001116757)	8E-60
Trypsin	JX477218	<i>Cyprinus carpio</i> (BAL04385)	2E-82
Complement C1r/s-A	JX477219	<i>Cyprinus carpio</i> (BAB17845)	4E-36
Cathepsin E	JX477220	<i>Danio rerio</i> (AAI65335)	2E-34
Cathepsin L, 1 a	JX477221	<i>Cyprinus carpio</i> (BAD08618)	7E-95
Cathepsin D	JX477222	<i>Ctenopharyngodon idella</i> (AAN62917)	1E-38
Cathepsin B	JX477223	<i>Cyprinus carpio</i> (BAE44111)	2E-40
Prothrombin-like	JX477224	<i>Oreochromis niloticus</i> (XP_003457819)	2E-25
60S ribosomal protein L40, Ubiquitin	JX477228	<i>Danio rerio</i> (NP_001032190)	8E-45
A-52 residue ribosomal protein fusion product 1			
FBR-musv ubiquitously expressed; ribosomal protein S30	JX477230	<i>Danio rerio</i> (NP_957031)	2E-45
Redox			
Selenoprotein P, plasma, 1a	JX477233	<i>Danio rerio</i> (NP_840082)	3E-54
Glutathione peroxidase 1a	JX477232	<i>Cyprinus carpio</i> (ADK26519)	2E-59
Glutathione peroxidase 3 (plasma)	JX477233	<i>Danio rerio</i> (NP_001131027)	6E-36
Glutathione peroxidase 4a	JX477234	<i>Carassius auratus</i> (ABO36294)	9E-67
Glutathione S-transferase theta 1a	JX477235	<i>Danio rerio</i> (XP_692427)	2E-151
Pi-class glutathione S-transferase	JX477237	<i>Carassius auratus</i> (ABF57553)	2E-41
Rho-class glutathione S-transferase	JX477236	<i>Cyprinus carpio</i> (ABD67511)	3E-79
Catalase	JX477239	<i>Ctenopharyngodon idella</i> (ACL99859)	7E-40
Ferritin, middle subunit-like	JX477240	<i>Danio rerio</i> (XP_687175)	5E-108
Natural killer cell enhancing factor B (NKEF-B)	JX477241	<i>Cyprinus carpio</i> (ABC59223)	4E-140
SOD (Cu/Zn)	JX477242	<i>Danio rerio</i> (NP_571369)	6E-57

Table 2 (continued)

Putative function	GenBank	Species (Accession no.)	E-value
Signal transduction			
Integrin, β 1b	JX477244	<i>Danio rerio</i> (AAI62706)	2E-46
Stress protein			
ATP synthase F0 subunit 6	JX477246	<i>Carassius auratus</i> (ABQ01626)	2E-46
Hsp5	JX477247	<i>Salmo salar</i> (NP_001135114)	7E-34
Hsp10	JX477248	<i>Danio rerio</i> (NP_571601)	5E-43
Tumor rejection antigen (gp96)	JX477250	<i>Danio rerio</i> (NP_937853)	7E-98
Acute phase response/inflammation			
Complement component C3	JX477251	<i>Cyprinus carpio</i> (BAA36619)	4E-66
Complement component C4-1	JX477252	<i>Cyprinus carpio</i> (BAB03284)	4E-35
Complement component C4-2	JX477253	<i>Cyprinus carpio</i> (BAB03285)	2E-63
Complement component C7	JX477254	<i>Hypophthalmichthys molitrix</i> (AFH01333)	1E-52
Complement component C9	JX477255	<i>Ctenopharyngodon idella</i> (ABN49522)	1E-47
Complement B/C2-A2	JX477256	<i>Cyprinus carpio</i> (BAA78416)	7E-65
Complement control protein factor I-B,	JX477257	<i>Cyprinus carpio</i> (BAB88921)	1E-75
Complement factor H like 4	JX477258	<i>Danio rerio</i> (CAP09610)	7E-31
Heme oxygenase	JX477259	<i>Ictalurus furcatus</i> (ADO27993)	4E-77
Macrophage stimulating 1	JX477260	<i>Danio rerio</i> (AAH84690)	1E-45
Transferrin variant B	JX477264	<i>Carassius gibelio</i> (AAL57602)	3E-30
Transferrin variant C	JX477262	<i>Carassius gibelio</i> (AAL57603)	2E-120
Transferrin variant D	JX477263	<i>Carassius gibelio</i> (AAM90970)	4E-64
Transferrin variant E	JX477261	<i>Carassius gibelio</i> (AAM90971)	4E-131

Abbreviations: FBR-musv, Finkel-Biskis-Reilly murine sarcoma virus; Hsp, heat shock protein; SOD, superoxide dismutase.

innate immune response and the adaptive immune system of animals with functionally divided two gene categories, inducible chemokines (inflammatory) and constitutive chemokines (homeostatic) [1,34]. Based on the number and location of conserved cysteine residues in the N-terminus, chemokines are classified into CCL, CXCL, CX3CL, and XCL subfamilies. However, an appropriate phylogenetic backbone or gene annotation criteria are still not available in the teleost chemokine subfamily as yet. Therefore, the identification of chemokines and a relevant study in fish may provide fundamental views concerning the potential role of each chemokine gene in the complexity of the immune system. Of the chemokine family, we identified one CC chemokine cDNA designated as *Ca-CCL25* gene in the crucian carp. The BLAST search and amino acid analysis showed that *Ca-CCL25* gene was close to zebrafish *CCL25s* gene (BAF98242) with 72% sequence similarity. As shown in Fig. 4, the *Ca-CCL25* transcript was detected strongly in kidney and spleen tissues. In head kidney, gill, and intestine tissues, moderate transcript levels of the *Ca-CCL25* gene were observed but only slight expressions were observed in other tissues. This result was similar from that reported previously in several fish such as *Pseudosciaena crocea* [59,61], *Cynoglossus semilaevis* [26], *Rachycentron canadum* [48], and *Oplegnathus fasciatus* [20] but different transcriptional profiles were also observed in *Oncorhynchus mykiss* [24] and *Miichthys miiuy* [10,56]. Thus, these results suggested that different types of CC chemokine genes would have evolved in different fish tissues. Also several reports supported this hypothesis. For example, Laing and Secombes [23] reported that both inducible and constitutive types of CC chemokines were presented in teleosts as different inducible patterns of 18 CC chemokines were observed upon recombinant TNF- α stimulation in the rainbow trout *O. mykiss*. Also, Borza et al. [4] suggested that CC chemokines would be regarded as either inducible or dual-function rather than a constitutive one based on tissue-specific different inducibilities of 12 CC chemokines of the Atlantic cod *Gadus morhua*. Particularly, high levels of different CC chemokine transcripts in kidney and spleen tissues were strongly correlated with the immune response against immune challenges in teleosts [29,59,61,8,26,48]. In teleosts, Trede et al. [50] suggested that kidney and spleen tissues were important immune-regulating organs for activation and interaction of chemokines as diverse lymphocytes, T and B cells, and myeloid

cells were also found in these tissues. Although *CCL25* is essential for colonization of T cells in thymus during early development of zebrafish [28], less is known in other teleosts. Therefore, the tissue-specific expression profile of *Ca-CCL25* gene will be helpful to understand its potential immune-relevant roles such as inflammatory and/or homeostatic functions in kidney and spleen tissues of the crucian carp.

3.3.3. Cathepsin

Cathepsins are lysosomal cysteine proteases for maintaining homeostasis and are among the most thoroughly studied hydrolases [3]. To date, over a dozen cathepsin proteases have been cloned and characterized in a number of mammalian and fish species (cathepsins B, C, F, H, K, L, O, S, T, U, V, W, and X). Cathepsins are classified into different subfamilies based on conserved domains/motifs and phylogenetic relationship with their substrates [22]. In the crucian carp, four cathepsin homologues (*Ca-CatB*, *Ca-CatD*, *Ca-CatE*, and *Ca-CatL*) were identified by pyrosequencing and *in silico* analysis. Based on overall transcriptional expressions of these genes in different tissues (Fig. 5), the constitutive expression of these cathepsins in immune and non-immune tissues indicates the pleiotropic nature of their functions and their crucial roles in homeostasis. Particularly, transcriptional abundance of *Ca-CatB* was measured in crucian carp, and revealed that the *Ca-CatB* transcript was mainly expressed in liver, intestine, and both kidney tissues. The *Ca-CatD* transcripts were distributed ubiquitously in most tissues except in eye, muscle, and skin. However, mRNA levels of *Ca-CatD* in kidney, liver, and spleen were greater than in other tissues. Study of expression of *Ca-CatE* at different tissues of the crucian carp indicated that the highest levels of transcripts were observed in kidney and liver. Heart, head kidney and intestine showed almost the same levels of mRNA expression. Tissue-specific differences in the transcriptional abundance of *Ca-CatL* were observed. In kidney, liver, and spleen, mRNA levels of *Ca-CatL* were highly detected, while in other tissues, the *CatL* transcripts were distributed ubiquitously in all the tissues but at a low level. Similarities and discrepancies of these four genes on tissue-specific transcriptional expressions have been reported in several fish such as *Paralichthys olivaceus* [59,61], *Cynoglossus semilaevis* [7], and *Oplegnathus fasciatus* [54] for *CatB* gene; *Oncorhynchus mykiss*

Table 3
Oligonucleotides used for real-time RT-PCR.

#	Gene	Oligonucleotide sequence	Tm (°C)	Length (bp)
1	Coagulation factor II	F: TTCCACCTCGGTCATCATC R: AATGCCAATCTGATACCAGC	56 55	148
2	Coagulation factor VII	F: TTGTAGAGGACATGATGGCAG R: GGTGTAATCCCATAGTTCCC	56 55	118
3	Coagulation factor X	F: CAGAAGTTAACGGTGCCATAC R: CCCTGACATGCATCTTTCTC	55 55	119
4	Coagulation factor V	F: TGGCCAGACTACATAATACAGG R: TGCTTGGCTCCTTGTGTG	55 55	124
5	C-type lectin	F: AGTGTGGAAATGGGTGGAC R: TGAATCACATGGCACGTC	54 55	158
6	Fibrinogen, β polypeptide	F: ATTCTATCAGCTGGTCTTACTC R: CTCAGGTATCCATTTGTCTG	55 55	151
7	Fibrinogen, γ polypeptide	F: GACTAGCCACATTCAAATC R: CGAAGTCGAAACCATCAAAG	55 56	106
8	Fibrinogen α chain	F: TGATAGGGACAGCGATAAGTG R: AATCTCATAAGGGACTTTGGTG	55 56	148
9	F11 receptor	F: TCTTCAATGGTAGCCTGGAG R: GCTGGTGTITGGCTGTTG	55 55	148
10	Interlectin 3	F: GAGATGGTACATGGGCAAAC R: GCTTATTATTGGGAACGTGC	55 55	132
11	CD59 glycoprotein	F: CCCTGAGATGTCAAACCTGTG R: ATGCACCTCCGGAAAGTAAAG	55 55	139
12	Integral membrane protein 2Bb	F: CATCATCACCCGAAACACATC R: GCTGTCCATCCTTCTGTAATC	55 56	139
13	Liver-expressed antimicrobial peptide 2B	F: ATACTTCATCGCAAGATCCG R: AGGCTCATTAACGAACAGTG	55 55	135
14	Programmed cell death 10	F: CTGCAGATGACGTGGAAGAG R: CCGTGTCCAGTAGTTCTTTTATG	56 57	183
15	Calmodulin	F: AAGATGAAGGACACGGACAG R: TCCTCATCTGTTAACTTCTCCC	54 55	137
16	Complement c1q, tumor necrosis factor-related protein	F: CAAGTACAGAGGATACAAACCG R: GCCTGAGTACCTGAAGTAATACAC	54 55	152
17	Myeloid cell leukemia sequence 1B	F: CGTTGTGAAGCACGAACTG R: GTGATGCCATCGCTGAAG	55 55	120
18	α -1-microglobulin	F: AGTGTGTTCCAGGTGAGCAG R: GGGACTTACACGCTTCAGG	55 55	138
19	α -2-macroglobulin	F: TCAACAACAGAATGAAGGGTG R: GGACTTACAACATGACAAACCTC	56 56	131
20	β -2-microglobulin	F: GTGCTGTACATCACTGTCCAAG R: TGATATCTGGAGGTGGAAG	55 54	130
21	MHC class I antigen	F: CAGTGGCCACAAGGAATCATC R: TCAGGCTGCTCTTCCATAC	56 55	135
22	Eukaryotic translation initiation factor 3 subunit M, dendritic cell protein	F: TCTTGCAATAGACCAGACACAG R: AGATTCTGTTCCAGGAGCAC	55 56	117
23	Leukocyte cell-derived chemotaxin 2	F: TACGAAGTATGGCTGTGGAAG R: AATGGAGCATAAACTGTGGC	56 55	108
24	Chemokine CCL-c25s	F: CCTGAGAAATGTTGCTGGTC R: TGTGATAAACCACAATGCCAG	55 55	111
25	Kininogen 1	F: ACCTGCAACAACCGGATAAG R: GACACTGGCGACTTTAGAG	55 55	129
26	Secreted immunoglobulin domain 4	F: CAGTCATCTCCTGGTCCAAG R: ATGTAAGTGGCTGAGTCTCTG	55 54	109
27	Carboxypeptidase B2	F: AGGAGGCTCAACAAAGATAC R: AAGCTCAAAGGTGAAGGAATAC	55 55	143
28	Signal sequence receptor, γ	F: CAAGCACAAAGTCCGCTCAG R: ATCCTCTCGTCTCTCTCTTTG	55 55	111
29	Fetuin-A, α -2-HS-glycoprotein	F: AGACTAGAACAGAAACACAGACG R: TGTGAAGCTGGTTCGGTG	56 55	116
30	Basic transcription factor 3	F: GAAATTAGCCAAGTTGCAGG R: GCCAGAGATGTTATTTACTCCC	55 55	148
31	Inter- α (globulin) inhibitor H3	F: CAGAGTCAAACCAACAGC R: TGATGAATGTACCGTATGCG	55 55	101
32	Inter- α -trypsin inhibitor heavy chain H4 precursor	F: AGTTGCTCGCAGGATTATG R: GCTGGTCTTGGTAAGGTTTG	56 55	133
33	Serine (or cysteine) proteinase inhibitor, clade C (antithrombin)	F: GACGATAAATGAATGGATAGCC R: TGACCTTTGAAGTAGATTGGC	56 56	120
34	Serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor)	F: AGTGAATGAAGAAGGTACCCGAG R: CCCATGAACACAACCGCAG	56 55	141
35	Serine (or cysteine) proteinase inhibitor, clade A (α -1 antiproteinase), member 1	F: TAGAGGGGAAGTGGGATAAAC R: TGGTTGACAGGGTCTTGATAG	55 55	138
36	Serpin peptidase inhibitor, clade A (α -1 antiproteinase), member 7	F: TCGAACTCACTTAAAGTCTCTC R: TCCACTTTTCAATGTGCTGTC	55 56	141
37	Serpin peptidase inhibitor, clade F, member 2B	F: AACTTGACTTCAACCCAGGACC R: CTGGACATATGCTGACACTGG	56 56	130
38	C1 inhibitor precursor	F: TTCTCTGCTCTGCTACCG	55	114

Table 3 (continued)

#	Gene	Oligonucleotide sequence	Tm (°C)	Length (bp)
39	Trypsin	R: ACTTCGAGCACCAATAGC F: CCGCTACCCTCAATCAGTAC	55 55	107
40	Complement C1r/s-	R: AGTGGAGCTCATGGTGTTC F: CTGTAATGGAGGGCAAGATG	54 56	139
41	Cathepsin E	R: TCAGTGACCAAATAATCCTCG F: CATCAACTGGATTCCAGTCAC	55 55	138
42	Cathepsin L, 1a	R: ATTAGAGAAGTCCCGTGTC F: GATGAATGGCTACAAACACAAG	55 55	127
43	Cathepsin D	R: ATCTTTCACGGGAGTCACG F: TGGGTGACTCTGTGTAAGG	55 55	147
44	Cathepsin B	R: TTACAGTCTACCAGAACTCTCC F: TGAGCATCATGTGAATGGC	55 55	122
45	Prothrombin-like	R: TTCCAAAGTGTTCCTCC F: ATAAAGAGAAAGAGTTGCTGC	55 56	116
46	60S ribosomal protein L40,	R: GTAATGTTCCGCCTGCTC F: CTACAACATTCAGAAAGAGTCCAC	56 56	123
47	FBR-musv ubiquitously expressed; ribosomal protein S30	R: CACTTGCCGAGATCATC F: TGTCTCTCTGGACTGTGGTG	55 55	139
48	Selenoprotein P, plasma, 1a	R: TTCTCCTGTTTCCACTTTG F: TGAATATCACCTCTACGCC	55 55	125
49	Glutathione peroxidase 1a	R: GGAAGAGACAGATGGTAAGTG F: GGATTACACCCAGATGAACG	56 55	128
50	Glutathione peroxidase 3 (plasma)	R: TCAGAGACAGCAGAAATTCATC F: CACTGCAACAGGAGTTACGAG	54 56	128
51	Glutathione peroxidase 4a	R: ATTGCCCTGGACGGACATAC F: TTGAAATCAGGAACCCAGG	55 55	98
52	Glutathione S-transferase theta 1a	R: GCCATTACATCAATCTTACTG F: TGGAAGTGTATCTCGAATTCG	55 55	126
53	Pi-class glutathione S-transferase	R: ATTCCTCTCCATATTGTTCCACC F: ACCTTCTGCTGAATCATAAAGTC	55 55	124
54	Rho-class glutathione S-transferase	R: AAGTTTTCACACTCCAGGAGG F: AAGGAGAGACATGAATCAGC	56 55	123
55	Catalase	R: TGAAGTCTTACCAGCGAGG F: CTTTACAATGCCATCTCCAAC	55 55	140
56	Ferritin, middle subunit-like	R: ATCAGAGGAAAGTCTTTATGGG F: AGACCCTCATCTGTGACTTC	55 55	114
57	Natural killer cell enhancing factor B (NKEF-B)	R: TTGTTGCCAGCATCCATC F: TCTGTAGACGAGACCTGAGAC	55 56	112
58	SOD (Cu/Zn)	R: GCACATCTGGAACAATAGTGTG F: CGCACTACAACCCTCATAATC	55 55	110
59	Integrin, β 1b	R: GTCAATTTCTGCAACACCATC F: ATCTGCCCAACTGTCTCTG	55 55	143
60	ATP synthase F0 subunit 6	R: AGTTCTCCTTCTTCTCACC F: ATCGGAATCGCAAATCAAC	54 56	124
61	Hsp 5	R: GGGCTAATGGTCGAATAAGTAG F: GACCTGAAGAAGACCGACATC	56 56	119
62	Hsp 10	R: TCCGGTTGATTCTCTG F: AAAGTCTCAAGCAAAGTGC	56 55	119
63	Tumor rejection antigen (gp96)	R: CTGGCAGCAGAACTTTATCC F: AGTTCGGCACCAACATTAAG	55 55	140
64	Complement component C3	R: CTCTCTTCTATCTCTCCAC F: TGAAACAGGACTCGGACATC	55 55	147
65	Complement component C4-1	R: GGTACGACTTGCTTCCAC F: AACGACGGACAACATTTAG	55 56	111
66	Complement component C4-2	R: AAGAAGATGGCAGGCAGC F: GTGTCAATGTGCAGAAAGAGG	56 55	130
67	Complement component C7	R: ATCCACAATGACTTCAAATGC F: CTCGTTTGTGTAATAATCTG	55 55	133
68	Complement component C9	R: GAAGGATCAAAGCAGTGGG F: GAGTGAAATGCATGTGCCTAC	55 55	103
69	Complement B/C2-A2	R: CCCAGTTTCCCAGTTGTTC F: TTGGAGTTGGAGATGAGGTG	55 55	143
70	Complement control protein factor I-B,	R: TACTCTGGTCTCATCAATC F: GACATCGCTTGGTGCAG	55 55	127
71	Complement factor H like 4	R: TGGTGCAGGTGTCATTGG F: CGAAGTGGACACAGGTGAC	55 56	123
72	Heme oxygenase	R: TGCCTCAACACATTTCCG F: GAGTATCTGGTGGCTCATGC	55 55	116
73	Macrophage stimulating 1	R: AACACAAGCCCTCTCCG F: TACACCCTGACCCCTGAACC	55 55	130
74	Transferrin variant B	R: GACACTCTTTCCCGCTCAC F: GAGAAGATGCACGGAAAGAC	55 54	134
75	Transferrin variant C	R: CTCCTGAAGGCATTTAGTGG F: GTTATGATGAAATCCACCG	55 55	136
		R: TCCTTGGCCGCTGATGAC	55	

(continued on next page)

Table 3 (continued)

#	Gene	Oligonucleotide sequence	Tm (°C)	Length (bp)
76	Transferrin variant D	F: TATTATGGCTATGATGGGGC R: CTCAGATCCTTAGCCCACTC	55 56	126
77	Transferrin variant E	F: AGGGCATCATCTGTGGAAG R: TTGCTCAACCATAACTGGAAC	55 55	126




Fig. 2. Amplicon identity of entire immune-relevant genes annotated in the EST library from *C. auratus*. Gene name for each number was incorporated in Table 3. A 100 bp ladder marker was used for each load.

[6], *Danio rerio* [39], *Scophthalmus maximus* [19], *Paralichthys olivaceus* [33], *Ictalurus punctatus* [16], and *Ctenopharyngodon idella* [13] for *CatD* gene; *Ictalurus punctatus* [57] and *Oplegnathus fasciatus* [54] for *CatL* gene, respectively, indicating that the gene regulation of cathepsins is conserved with some divergence in fish.




Fig. 3. Tissue-specific mRNA expressions of *Ca-CTL* gene in different tissues of the crucian carp. *C. auratus* β -actin gene was used as a reference housekeeping gene to normalize the expression. Each value represents an average of three replicate samples (\pm S.D.). Significant difference over the lowest value are indicated by different small letters on the data bars ($P < 0.05$) analyzed by multiple-comparison ANOVA.

In vertebrate tissues, cathepsins are known to be ubiquitously expressed with different and specific roles in almost all cell types of the crucian carp as well. Also, our results suggest that the cathepsin family of the crucian carp is involved in a number of physiological processes that have a potent role in host defense through innate immunity. Due to the versatility of expression of the cathepsin family, further studies are needed to unveil their functions in each tissue of crucian carp.

3.3.4. Natural killer cell enhancing factor

Natural killer (NK) cell enhancing factor (NKEF) belongs to the newly defined peroxiredoxin (Prx) family. Originally, NKEF as a cytosolic protein of the human red blood cells has been known to enhance NK cytotoxic activity against the K562 tumor cell line [44]. In mammals, NKEF protein is consisted of two highly homologous genes as Prx1 (NKEF-A) and Prx2 (NKEF-B). Recently, NKEFs have been cloned from diverse teleosts, and their susceptibility against immune challenges suggested a role for NKEFs in inflammation as an important regulator in antitumoral and antiviral defense (summarized in a recent publication; [15]). In the crucian carp, *NKEF-B* gene (*Ca-NKEF-B*) was annotated from EST database, and its tissue-preferential transcript expression was analyzed. As shown in Fig. 6, transcriptional abundance of *Ca-NKEF-B* was ubiquitously detected in almost tissues, while high levels were measured in kidney, heart, and head kidney tissues. This kind of expression




Fig. 4. Tissue-specific mRNA expressions of *Ca-CCL25* gene in different tissues of the crucian carp. *C. auratus* β -actin gene was used as a reference housekeeping gene to normalize the expression. Each value represents an average of three replicate samples (\pm S.D.). Significant difference over the lowest value are indicated by different small letters on the data bars ($P < 0.05$) analyzed by multiple-comparison ANOVA.




Fig. 6. Tissue-specific mRNA expression of *NKEF* gene in different tissues of the crucian carp. *C. auratus* β -actin gene was used as a reference housekeeping gene to normalize the expression. Each value represents an average of three replicate samples (\pm S.D.). Significant difference over the lowest value are indicated by different small letters on the data bars ($P < 0.05$) analyzed by multiple-comparison ANOVA.

patterns were observed in different tissues of diverse teleosts but its tissue-specific function is not clear as yet. Possibly, ubiquitous profiles can be explained as NKEFs are probably associated with forming tissue [43]. In fact, peroxiredoxins are important proteins in the antioxidative defense system and have relevant functions in immunity [38]. Marionnet et al. [30] reported that several immunostimulants (lipopolysaccharide, dextran sulfate, and tilorone) could induce intracellular ROS production by macrophages in the head kidney and the spleen tissues of common carp. Thus, we suggest that *Ca-NKEF-B* might have an antioxidant role in diverse tissues to prevent immune challenges-induced oxidative stress and further damage, resulting in ubiquitous transcript expression in the crucian carp.

3.3.5. Complement component

Complement system is a part of the immune defense system and is regulated by complement control proteins. Activation of the complement system as a supportive defense line of innate

immunity triggers the assembly of the terminal complement complexes against immune challenges, resulting in the induction of the characteristic complement-mediated cytolysis [41]. In the crucian carp, five complement components, *Ca-C3*, *Ca-C4-1*, *Ca-C4-2*, *Ca-C7*, and *Ca-C9*, were annotated from the EST database. The complement system is composed of three major pathways such as the classical complement pathway (CCP), the alternative complement pathway (ACP), and the lectin complement pathway (LCP) [5]. Of complements, the complement component 3 (C3) is a central protein of the entire complement system. Upon immune challenges, C3 induces an amphipathic pore structure as a membrane attack complex (MAC) that is composed by C5b, C6, C7, C8, and C9 molecules [40]. C4 molecule is known to be involved in the prevention of the autoimmune disease with maintenance of the B-cell tolerance in mammal [9]. Also, C4 is a key molecule for the activation of both the CCP and the LCP in teleosts [18]. These complement components were highly detected in liver tissues of the crucian carp (Fig. 7). Our results strongly supported the common




Fig. 5. Tissue-specific mRNA expressions of four cathepsin genes in different tissues of the crucian carp. *C. auratus* β -actin gene was used as a reference housekeeping gene to normalize the expression. Each value represents an average of three replicate samples (\pm S.D.). In each gene, significant differences in different tissues over the lowest value are indicated by different small letters on the data bars ($P < 0.05$) analyzed by multiple-comparison ANOVA. In each tissue, significant difference between different transcripts is indicated by capital letters on the data bars ($P < 0.05$) analyzed by one-way ANOVA followed by Tukey's test. When none of the characters between data bars match, values are considered statistically insignificant.




Fig. 7. Tissue-specific mRNA expressions of five complement components in different tissues of the crucian carp. *C. auratus* β -actin gene was used as a reference housekeeping gene to normalize the expression. Each value represents an average of three replicate samples (\pm S.D). In each gene, significant differences in different tissues over the lowest value are indicated by different small letters on the data bars ($P < 0.05$) analyzed by multiple-comparison ANOVA. In each tissue, significant difference between different transcripts is indicated by capital letters on the data bars ($P < 0.05$) analyzed by one-way ANOVA followed by Tukey's test. When none of the characters between data bars match, values are considered statistically insignificant.

feature that complement components are primarily synthesized as inactive precursors in hepatocytes. Also, transcriptional expression of the complement components in other tissues of the crucian carp can be explained by extrahepatic synthesis of several complement proteins, as different tissues are known to be associated with complement production for further inflammatory responses [25,32]. In most complement components identified in teleosts, a vast majority of transcriptional abundance was observed in liver tissues but only slightly expressed in other tissues. Thus, we can conclude that liver of the crucian carp synthesizes complement components with several extrahepatic sources.

4. Conclusion

Using the pyrosequencing method, we confirmed putative immune-related gene information in the crucian carp, *C. auratus* as studied in numerous previous reports, and demonstrated the effectiveness of the gene cloning method with pyrosequencing. Immune-related gene information of the crucian carp provides clues to develop immune response analysis in this species, as the crucian carp has been known to be one of most important commercial species in Asian aquaculture. Ongoing studies are targeting the specific role of the entire set of immune-relevant genes upon immune challenge in this species.

Acknowledgments

We thank Dr. Hans-U. Dahms for his comments on the manuscript. This research was a part of the project titled "Korea-Polar Ocean Development: K-POD" (project no. PM12030) funded by the Ministry of Oceans and Fisheries, Korea.

References

- [1] Alejo A, Tafalla C. Chemokines in teleost fish species. *Dev Comp Immunol* 2011;35:1215–22.
- [2] Aoki T, Takano T, Santos MD, Kondo H, Hirono I. Molecular innate immunity in teleost fish: review and future perspectives. *Fisheries for global welfare and environment*. In: 5th world fisheries congress 2008. pp. 263–76.
- [3] Brix K, Dunkhorst A, Mayer K, Jordans S. Cysteine cathepsins: cellular roadmap to different functions. *Biochimie* 2008;90:194–207.
- [4] Borza T, Stone C, Rise ML, Bowman S, Johnson SC. Atlantic cod (*Gadus morhua*) CC chemokines: diversity and expression analysis. *Dev Comp Immunol* 2010;34:904–13.
- [5] Boshra H, Li J, Sunyer JO. Recent advances on the complement system of teleost fish. *Fish Shellfish Immunol* 2006;20:239–62.
- [6] Brooks S, Tyler CR, Carnevali O, Coward K, Sumpter JP. Molecular characterization of ovarian cathepsin D in the rainbow trout, *Oncorhynchus mykiss*. *Gene* 1997;201:45–54.
- [7] Chen L, Sun L. Cathepsin B of *Cynoglossus semilaevis*: identification, expression, and activity analysis. *Comp Biochem Physiol B* 2012;161:54–9.
- [8] Chen SL, Liu Y, Dong XL, Meng L. Cloning, characterization, and expression analysis of a CC chemokine gene from turbot (*Scophthalmus maximus*). *Fish Physiol Biochem* 2010;36:147–55.
- [9] Chen Z, Koralov SB, Kelsø G. Complement C4 inhibits systemic autoimmunity through a mechanism independent of complement receptors CR1 and CR2. *J Exp Med* 2000;192:1339–52.
- [10] Cheng YZ, Wang RX, Sun YN, Xu TJ. Molecular characterization of miiuy croaker CC chemokine gene and its expression following *Vibrio anguillarum* injection. *Fish Shellfish Immunol* 2011;31:148–54.
- [11] Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 2005;21:3674–6.
- [12] Day AJ. The C-type carbohydrate recognition domain (CRD) superfamily. *Biochem Soc Trans* 1994;22:83–8.
- [13] Dong ZD, Zhang J, Ji XS, Zhou FN, Fu Y, Chen W, et al. Molecular cloning, characterization and expression of cathepsin D from grass carp (*Ctenopharyngodon idella*). *Fish Shellfish Immunol* 2012;33:1207–1204.
- [14] Encinas P, Rodriguez-Milla MA, Novoa B, Estepa A, Figueras A, Coll J. Zebrafish fin immune responses during high mortality infections with viral haemorrhagic septicaemia rhabdovirus. A proteomic and transcriptomic approach. *BMC Genomics* 2010;11:518.
- [15] Esteban MA, Chaves-Pozo E, Arizcun M, Meseguer J, Cuesta A. Regulation of natural killer enhancing factor (*NKEF*) genes in teleost fish, gilthead seabream and European sea bass. *Mol Immunol* 2013;55:275–82.
- [16] Feng T, Zhang H, Liu H, Zhou Z, Niu D, Wong L, et al. Molecular characterization and expression analysis of the channel catfish cathepsin D genes. *Fish Shellfish Immunol* 2011;31:164–9.
- [17] Gardner JP, Durso RJ, Arrigale RR, Donovan GP, Maddon PJ, Dragic T, et al. L-SIGN (CD 209L) is a liver-specific capture receptor for hepatitis C virus. *Proc Natl Acad Sci U S A* 2003;100:4498–503.
- [18] Gonzalez SF, Buchmann K, Nielsen ME. Complement expression in common carp (*Cyprinus carpio* L.) during infection with *Ichthyophthirius multifiliis*. *Dev Comp Immunol* 2007;31:576–86.
- [19] Jia A, Zhang XH. Molecular cloning, characterization and expression analysis of cathepsin D gene from turbot *Scophthalmus maximus*. *Fish Shellfish Immunol* 2009;26:606–13.
- [20] Kim J-W, Kim E-G, Kim D-H, Shim SH, Park C-I. Molecular identification and expression analysis of the CC chemokine gene in rock bream (*Oplegnathus*

- fasciatus*) and the biological activity of the recombinant protein. *Fish Shellfish Immunol* 2013;34:892–901.
- [21] Kondo H, Yeu Tzeh AG, Hirono I, Aoki T. Identification of a novel C-type lectin gene in Japanese flounder, *Paralichthys olivaceus*. *Fish Shellfish Immunol* 2007;23:1089–94.
- [22] Lacaille F, Kaleta J, Brömme D. Human and parasitic papain-like cysteine proteases: their role in physiology and pathology and recent developments in inhibitor design. *Chem Rev* 2002;102:4459–88.
- [23] Laing KJ, Secombes CJ. Trout CC chemokines: comparison of their sequences and expression patterns. *Mol Immunol* 2004;41:793–808.
- [24] Lally J, Al-Anouti F, Bols N, Dixon B. The functional characterisation of CK-1, a putative CC chemokine from rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol* 2003;15:411–24.
- [25] Laufer J, Katz Y, Passwell JH. Extra-hepatic synthesis of complement proteins in inflammation. *Mol Immunol* 2001;38:221–9.
- [26] Li YX, Sun JS, Sun L. An inflammatory CC chemokine of *Cynoglossus semilaevis* is involved in the immune defense against bacterial infection. *Fish Shellfish Immunol* 2011;31:446–52.
- [27] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real time quantitative PCR and the $2^{-\Delta\Delta C_t}$ method. *Methods* 2001;25:402–8.
- [28] Lu IN, Chiang BL, Lou KL, Huang PT, Yao CC, Wang JS, et al. Cloning, expression and characterization of CCL21 and CCL25 chemokines in zebrafish. *Dev Comp Immunol* 2012;38:203–14.
- [29] Mackenzie S, Liarte C, Ilied V, Planas JV, Tort L, Goetz FW. Characterization of a highly inducible novel CC chemokine from differentiated rainbow trout (*Oncorhynchus mykiss*) macrophages. *Immunogenetics* 2004;56:611–5.
- [30] Marionnet D, Deschaux P, Reynaud S. Possible implication of macrophages in the regulation of cytochrome P450 activities in carp (*Cyprinus carpio*). *Fish Shellfish Immunol* 2006;21:80–91.
- [31] Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, et al. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 2005;437:376–80.
- [32] Morgan B, Gasque P. Extrahepatic complement biosynthesis: where, when and why? *Clin Exp Immunol* 1997;107:1–7.
- [33] Park E-M, Kim Y-O, Nam B-H, Kong HJ, Kim W-J, Lee S-J, et al. Cloning and expression analysis of cathepsin D in the olive flounder *Paralichthys olivaceus*. *Biosci Biotechnol Biochem* 2009;73:1856–9.
- [34] Peatman E, Liu Z. Evolution of CC chemokines in teleost fish: a case study in gene duplication and implications for immune diversity. *Immunogenetics* 2007;59:613–23.
- [35] Pöhlmann S, Zhang J, Baribaud F, Chen Z, Leslie GJ, Lin G, et al. Hepatitis C virus glycoproteins interact with DC-SIGN and DC-SIGNR. *J Virol* 2003;77:4070–80.
- [36] Rhee J-S, Kim B-M, Kim R-O, Choi B-S, Choi I-Y, Lee Y-M, et al. Analysis of expressed sequence tags from the liver and ovary of the euryhaline hermaphroditic fish, *Kryptolebias marmoratus*. *Comp Biochem Physiol D* 2011;6:244–55.
- [37] Rhee J-S, Kim B-M, Jeong C-B, Horiguchi T, Lee Y-M, Kim I-C, et al. Immune gene mining by pyrosequencing in the rockshell, *Thais clavigera*. *Fish Shellfish Immunol* 2012;32:700–10.
- [38] Rhee SG, Chae HZ, Kim K. Peroxiredoxins: a historical overview and speculative preview of novel mechanisms and emerging concepts in cell signaling. *Free Radic Biol Med* 2005;38:1543–52.
- [39] Riggio M, Scudiero R, Filosa S, Parisi E. Sex- and tissue-specific expression of aspartic proteinases in *Danio rerio* (zebrafish). *Gene* 2000;260:67–75.
- [40] Rutkowski MJ, Sughrue ME, Kane AJ, Ahn BJ, Fang S, Parsa AT. The complement cascade as a mediator of tissue growth and regeneration. *Inflamm Res* 2010;59:897–905.
- [41] Sarma JV, Ward PA. The complement system. *Cell Tissue Res* 2011;343:227–35.
- [42] Savan R, Endo M, Sakai M. Characterization of a new C-type lectin from common carp *Cyprinus carpio*. *Mol Immunol* 2004;41:891–9.
- [43] Sauri H, Ashjian PH, Kim AT, Shau H. Recombinant natural killer enhancing factor augments natural killer cytotoxicity. *J Leukoc Biol* 1996;59:925–31.
- [44] Shau H, Gupta RK, Golub SH. Identification of a natural killer enhancing factor (NKEF) from human erythroid cells. *Cell Immunol* 1993;147:1–11.
- [45] Shi J, Zhang YB, Liu TK, Sun F, Gui JF. Subcellular localization and functional characterization of a fish *IRF9* from crucian carp *Carassius auratus*. *Fish Shellfish Immunol* 2013;33:258–66.
- [46] Somamoto T, Yoshiura Y, Nakanishi T, Ototake M. Molecular cloning and characterization of two types of *CD8 α* from ginbuna crucian carp, *Carassius auratus langsdorffii*. *Dev Comp Immunol* 2005;29:693–702.
- [47] Star B, Nederbragt AJ, Jentoft S, Grimholt U, Malmstrøm M, Gregers TF, et al. The genome sequence of Atlantic cod reveals a unique immune system. *Nature* 2011;477:207–10.
- [48] Su Y, Guo Z, Xu L, Jiang J, Wang J, Feng J. Identification of a cobia (*Rachycentron canadum*) CC chemokine gene and its involvement in the inflammatory response. *Fish Shellfish Immunol* 2012;32:204–10.
- [49] Takizawa F, Mizunaga Y, Araki K, Moritomo T, Ototake M, Nakanishi T. *GATA3* mRNA in ginbuna crucian carp (*Carassius auratus langsdorffii*): cDNA cloning, splice variants and expression analysis. *Dev Comp Immunol* 2008;32:898–907.
- [50] Trede NS, Langenau DM, Traver D, Look AT, Zon LI. The use of zebrafish to understand immunity. *Immunity* 2004;20:367–79.
- [51] Turk V, Stoka V, Vasiljeva O, Renko M, Sun T, Turk B, et al. Cysteine cathepsins: from structure, function and regulation to new frontiers. *Biochim Biophys Acta* 2012;1824:68–88.
- [52] Urabe S, Somamoto T, Sameshima S, Unoki-Kato Y, Nakanishi T, Nakao M. Molecular characterization of MHC class I and β -2 microglobulin in a clonal strain of ginbuna crucian carp, *Carassius auratus langsdorffii*. *Fish Shellfish Immunol* 2011;31:469–74.
- [53] Wei J, Xu D, Zhou J, Cui H, Yan Y, Ouyang Z, et al. Molecular cloning, characterization and expression analysis of a C-type lectin (*Ec-CTL*) in orange-spotted grouper, *Epinephelus coioides*. *Fish Shellfish Immunol* 2010;28:178–86.
- [54] Whang I, De Zoysa M, Nikapitiya C, Lee Y, Kim Y, Lee S, et al. Molecular characterization and expression analysis of Cathepsin B and L cysteine proteases from rock bream (*Oplegnathus fasciatus*). *Fish Shellfish Immunol* 2011;30:763–72.
- [55] Xiang LX, He D, Dong WR, Zhang YW, Shao JZ. Deep sequencing-based transcriptome profiling analysis of bacteria-challenged *Lateolabrax japonicus* reveals insight into the immune-relevant genes in marine fish. *BMC Genomics* 2010;11:472.
- [56] Xu TJ, Cheng YZ, Ge S, Wang RX. Molecular cloning, characterization, and expression analysis of a disease resistance related CC chemokine gene in miiuy croaker (*Miichthys miiuy*). *Aquaculture* 2011;318:25–32.
- [57] Yeh HY, Klesius PH. Channel catfish, *Ictalurus punctatus*, cysteine proteinases: cloning, characterisation and expression of cathepsin H and L. *Fish Shellfish Immunol* 2009;26:332–8.
- [58] Zelensky AN, Gready JE. The C-type lectin-like domain superfamily. *FEBS J* 2005;272:6179–217.
- [59] Zhang FT, Zhang YB, Chen YD, Zhu R, Dong CW, Li YY, et al. Expressional induction of *Paralichthys olivaceus* cathepsin B gene in response to virus, poly I: C and lipopolysaccharide. *Fish Shellfish Immunol* 2008;25:542–9.
- [60] Zhang H, Robison B, Thorgaard GH, Ristow SS. Cloning, mapping and genomic organization of a fish C-type lectin gene from homozygous clones of rainbow trout (*Oncorhynchus mykiss*). *Biochim Biophys Acta* 2000;1494:14–22.
- [61] Zhang J, Ao J, Chen X. Molecular and functional characterization of a novel CC chemokine in large yellow croaker (*Pseudosciaena crocea*). *Fish Shellfish Immunol* 2008;25:664–71.
- [62] Zhang M, Hu YH, Sun L. Identification and molecular analysis of a novel C-type lectin from *Scophthalmus maximus*. *Fish Shellfish Immunol* 2010;29:82–8.
- [63] Zhang YB, Hu CY, Zhang J, Huang GP, Wei LH, Zhang QY, et al. Molecular cloning and characterization of crucian carp (*Carassius auratus* L.) interferon regulatory factor 7. *Fish Shellfish Immunol* 2003;15:453–66.
- [64] Zhang YB, Wang YL, Gui JF. Identification and characterization of two homologues of interferon-stimulated gene ISG15 in crucian carp. *Fish Shellfish Immunol* 2007;23:52–61.
- [65] Zlotnik A, Yoshie O, Nomiya H. The chemokine and chemokine receptor superfamilies and their molecular evolution. *Genome Biol* 2006;7:243.