

Effects of wind and sea ice on the seasonal variation of warm circumpolar deep water in the Amundsen Sea

Tae-Wan Kim¹, Ho Kyung Ha^{1,2}, Anna K. Wåhlin³, Jae Hak Lee⁴, Hyun Jung Lee¹ and SangHoon Lee¹

¹Korea Polar Research Institute, Incheon 406-840, South Korea

²Department of Ocean Sciences, Inha University, Incheon 402-751, South Korea


³Department of Earth Sciences, University of Gothenburg, Gothenburg, Sweden

⁴Korea Institute of Ocean Science and Technology, Ansan 425-600, South Korea

Introduction

West Antarctic ice sheet has experienced a pronounced mass loss in recent decades. The most rapidly changing region of the West Antarctic is the Amundsen Sea, where the intrusion of relatively warm circumpolar deep water (CDW) onto the continental shelf may help reduce ice thickness. Recently, the temporal variation in the flow of CDW was examined using a mooring current meter in the center of the deep inflow in the Dotson Trough. The existence of seasonal variation of bottom temperature and thickness of warm layer of CDW were measured from this mooring. Also, the velocity has strong barotropic fluctuations that correlate with the eastward wind at the shelf break on short-term scales. However, the bottom temperature could be shown to correlate with the wind from the time series. The main objective of this study was to examine the combined effect of wind and sea ice drift on the on-shelf flow of CDW in the Amundsen Sea. In order to understand the effects of wind and sea ice on the seasonal variation in the thickness of CDW, we calculate the ocean surface stress curl and Ekman pumping velocity in Amundsen Sea.

Materials and methods




Two oceanographic surveys were conducted by the IBRV Araon from 21 December 2010 to 23 January 2011, and from 31 January to 20 March 2012 (Figure 1). A total of 30 and 52 CTD stations were occupied during the surveys in 2011 and 2012, respectively. The temporal variability and properties of CDW were observed from a mooring (72.46°S, 116.35°W) in the eastern side of Dotson Trough (Figure 1) from 15 February 2010 to 1 March 2012. In order to calculate the Ekman pumping velocity, we use the reanalysis wind data and observed sea ice concentration and velocity data from satellite. Wind data were obtained from the ERA interim reanalysis data. Sea ice concentration data were obtained from the Nimbus-7 SMMR, DMSP, SSM/I, SSMIS. The sea ice velocity data were obtained from the Polar Pathfinder Daily 25 km EASE-Grid Sea Ice Motion Vectors Version 2 from 1990 to 2011.

Seasonal variation of the CDW layer


The thickness of the warm layer (identified here as $T > 0^\circ\text{C}$ isotherm) and the bottom temperature vary distinctly with season; both attain maxima in austral summer and minima in austral winter. The difference between the seasonal maximum and minimum in thickness is approximately 60-100m and the difference between maximum and minimum bottom temperature is approximately 1°C ($0.25\text{-}1.2^\circ\text{C}$). The time-series of the heat content mirrors the seasonal variation in the thickness of the MCDW layer.

Time-series variation of potential temperature, salinity, neutral density and depth averaged heat content at the mooring station. The white line in upper panel indicates the upper boundary of the warm layer (0°C isotherm).





Spatial and temporal variation of the ocean surface stress

To examine the effects of the varying Ekman pumping velocity on the seasonality of both the heat content and the position of the 0°C isotherm, a time series of the wind field and the sea ice extent were used. Maximum eastward ocean surface stress (0.5 dyne cm^{-2}) was at 66°S , where the sea ice concentration was less than 0.25. The minimum absolute value of ocean surface stress ($0.03 \text{ dyne cm}^{-2}$) was noted at the area of highest sea ice concentration (70°S). Stress curl and Ekman pumping velocity at the sea surface were calculated from the horizontal variation of ocean surface stress. An extensive area of positive wind stress curl and high Ekman pumping velocities ($>0.26 \text{ m day}^{-1}$) was located around 66°S . Between 68°S and 71°S , where sea ice concentration is higher than 40%, the stress curl is less than $0.5 \times 10^{-8} \text{ dyne cm}^{-3}$. Wind stress curl and sea ice concentration, both of which show pronounced seasonal variation, give rise to spatial variation in the Ekman pumping velocity. From February to April (FMA), north of 70°S , Ekman pumping exceeds 0.2 m day^{-1} due to latitudinal variation in the sea ice concentration and the wind. North of 68°S , despite the latitudinal variation of wind, Ekman pumping velocity decreases due to the horizontally homogenous sea ice concentration. A strong Ekman pumping occurs near the Amundsen Sea shelf break. From August to October (ASO), the sea ice zone extends north of the shelf break and the Ekman pumping velocity at the shelf break is notably weak during this season.



Discussion and conclusions

Both heat content and the position of the 0°C isotherm showed seasonal variation, attaining a maximum during austral summer and a minimum during austral winter. The meridional variation in the observed isohalines and isotherms north of Amundsen Sea shelf break suggests that the thermocline depth depended mainly on the upwelling/downwelling caused by local wind patterns and sea ice conditions. Therefore, it appears that the seasonal variation of the thickness of the CDW at the Amundsen Sea shelf break is mainly affected by the variation in local wind patterns and sea ice conditions. The monthly averaged sea ice concentration was less than 50% from January to April and higher than 50% for the rest of the year. The Ekman pumping velocity reached a maximum of 0.25 m day^{-1} in April when the sea ice concentration was less than 40% and the marginal ice zone was located at Amundsen Sea shelf break. In the austral winter, the sea ice concentration was high ($> 80\%$) and latitudinally homogenous at the Amundsen Sea shelf break; so the Ekman pumping was small, even though the eastward wind was stronger.



During the austral summer, the sea ice zone contracts because of atmospheric heating, and Ekman pumping occurs north of the sea ice boundary due to the latitudinal change in velocity of the eastward wind. Ekman pumping is especially strong around the Amundsen Sea shelf break or sea ice marginal zone due to the strong latitudinal gradient in surface stress (caused by sea ice). In contrast, during the austral winter, the sea ice zone extends further north (to about 65°S). Because of the horizontal homogeneity in sea ice concentration, there is less of a latitudinal gradient in surface stress across the sea ice covered areas of the Amundsen Sea shelf. Consequently, Ekman pumping velocity is also lower, although the latitudinal change in wind velocity in austral winter is greater than that in austral summer. Around the sea ice boundary, however, there is a rather large latitudinal difference in wind stress, due to the strong eastward wind speed and the steep latitudinal change in sea ice concentration.