

## Detection of Plutonium-239 in the Antarctic snow by inductively coupled plasma-sector field mass spectrometry

Plutonium is a useful time marker for dating ice cores and snow pits because it in the environment mainly originates from atmospheric nuclear weapons tests carried out since the 1950s. To determine  $^{239}\text{Pu}$  in snow pit samples, collected every 5 cm down to 4 m, covering ~50 years (1957-2007), at Dome Fuji in East Antarctica, we used an inductively coupled plasma-sector field mass spectrometer (ICP-SFMS) coupled to a high-efficiency sample introduction system. The main advantages of ICP-SFMS technique are rapidity of analysis and simple sample preparation method for  $^{239}\text{Pu}$  at femtogram levels in small-volume samples from snow/ice. However, this technique is prone to spectral interferences. The existence of high content of uranium in sample could lead to significant interferences with  $^{239}\text{Pu}$  owing to uranium hydride ( $^{238}\text{UH}^+$ ) formation. In this study, we found that the interference effect of  $^{238}\text{UH}^+$  was negligible when the  $^{238}\text{U}$  concentrations lower than  $10 \text{ pg g}^{-1}$ . In the snow pit samples, the  $^{238}\text{U}$  concentrations were lower than  $0.5 \text{ pg g}^{-1}$ . Accordingly,  $^{239}\text{Pu}$  signals were detected without  $^{238}\text{UH}^+$  interference. These suggest that this method can be widely used for the reconstruction of the fallout history of  $^{239}\text{Pu}$  and for the age constraint in other Antarctic Plateau sites.